


ISBN 978-81-180544-3-6



# ZOOLOGICAL TECHNIQUES: A Practical Guide for UG and PG

Mr. Kishor More ■ Ms. Sharvari Shah  
Dr. Kavita Kumbhar ■ Ms. Shaheen Shaikh



**OLYMPICK**  
PUBLISHER INDIA

<https://olympickpublisher.in/>

SEPTEMBER  
2026

---

# **ZOOLOGICAL TECHNIQUES: A PRACTICAL GUIDE FOR UNDERGRADUATE AND POSTGRADUATE STUDENTS**

Vol - I

2026

Mr. Kishor U. More · Ms. Sharvari V. Shah

Dr. Kavita Kumbhar · MS. Shaheen M. Shaikh



---

## ***Editors***

**Mr. Kishor U. More**

**Ms. Sharvari V. Shah**

**Dr. Kavita Kumbhar**

**MS. Shaheen M. Shaikh**

DOI : <https://doi.org/10.5281/zenodo.22762187>

ISBN 978-81-180544-3-6

<https://olympickpublisher.in/>



[contact@olympickpublisher.in](mailto:contact@olympickpublisher.in)

Registered Address: Olympick Publisher India 201, Shitole Empire Building,  
New Sangvi, Haveli, Pune - 411027

© The Author(s) and Editor(s) (if applicable) hold an exclusive license from Olympick Publisher India. Any reproduction, distribution or transmission of this publication in any form requires prior written authorization from the publisher except when brief quotations are used in academic research or critical reviews.

At publication time the publisher, authors / editors made every possible effort to provide information and advice in this book which is both accurate and current. The authors/ editors give no expressed or implied warranties about the material's completeness or correctness and they accept no responsibility for mistakes or missing information.

When disposing of this product, please ensure proper paper recycling.

---

## Contents

|                                                                            |    |
|----------------------------------------------------------------------------|----|
| <b>PART A · FOUNDATION TECHNIQUES</b>                                      | 1  |
| 1. Introduction to Zoological Techniques and Laboratory Safety             | 1  |
| 2. Collection, Preservation and Museum Techniques                          | 05 |
| 3. Microscopy: Principles and Applications                                 | 10 |
| <b>PART B · HISTOLOGICAL AND CYTOLOGICAL TECHNIQUES</b>                    |    |
| 4. Microtomy and Histological Techniques                                   | 14 |
| 5. Histochemistry and Immunohistochemistry                                 | 17 |
| 6. Chromosome Preparation and Karyotyping                                  | 21 |
| <b>PART C · BIOPHYSICAL AND BIOCHEMICAL TECHNIQUES</b>                     |    |
| 7. Centrifugation Techniques                                               | 25 |
| 8. Spectrophotometry and Colorimetry                                       | 28 |
| 9. Chromatographic Techniques                                              | 31 |
| 10. Electrophoretic Techniques                                             | 35 |
| 11. Radioisotope Techniques in Biology                                     | 39 |
| <b>PART D · MOLECULAR BIOLOGY AND BIOINFORMATICS</b>                       |    |
| 12. Nucleic Acid Extraction and PCR-based Techniques                       | 42 |
| 13. Recombinant DNA Technology and Gene Analysis                           | 47 |
| 14. Bioinformatics Tools in Zoology                                        | 50 |
| <b>PART E · IMMUNOLOGY, TOXICOLOGY AND CELL CULTURE</b>                    |    |
| 15. Immunological Techniques                                               | 53 |
| 16. Toxicological and Bioassay Techniques                                  | 57 |
| 17. Animal Tissue Culture Techniques                                       | 60 |
| <b>PART F · ECOLOGICAL, WILDLIFE AND BEHAVIOURAL TECHNIQUES</b>            |    |
| 18. Ecological Sampling Techniques                                         | 64 |
| 19. Wildlife Census and Population Estimation Techniques                   | 68 |
| 20. Remote Sensing, GIS and Animal Behaviour Study                         | 71 |
| <b>PART G · BIostatISTICS AND RESEARCH METHODOLOGY</b>                     |    |
| 21. Biostatistics for Zoological Data                                      | 75 |
| 22. Research Methodology, Thesis Writing, Ethics and Intellectual Property | 79 |
| <i>Appendix I · Glossary of Technical Terms</i>                            | 83 |





# CHAPTER 1. INTRODUCTION TO ZOOLOGICAL TECHNIQUES AND LABORATORY SAFETY

## PART A · FOUNDATION TECHNIQUES

*Core skills basic to field collection, preservation and observation*

**Learning focus:** *UG entry point; PG refresher*

### Scope of Zoological Technique

A zoological technique is a defined, repeatable procedure used to collect, preserve, examine or analyse animal material for the purpose of scientific study. The subject spans four broad domains: field methods for collection and observation, laboratory methods for structural and biochemical analysis, molecular methods for genetic and diagnostic study, and quantitative methods for interpreting the resulting data.

Competence in technique converts a biological question into verifiable evidence. A student unfamiliar with fixation chemistry damages tissue before it reaches the microscope; a researcher unfamiliar with sampling design collects data unable to answer the stated hypothesis. This book treats technique as a subject in its own right, distinct from the taxonomy, physiology or ecology to which it is later applied.

### Organization of Laboratory Work

Every zoological laboratory exercise follows a common sequence: preparation of materials, execution of the procedure, observation or measurement, recording of results. Deviation from this sequence, particularly omission of the recording step, is the most frequent source of unusable data among beginners.

### Categories of Laboratory Instrumentation

The instrumentation used in a zoology laboratory can be grouped into four broad categories on the basis of function. Observational instruments, comprising the compound and stereo microscope, extend the resolving power of the unaided eye and are treated in detail in Chapter 3. Preparative instruments, comprising the microtome, centrifuge and autoclave, convert raw biological material into a form suitable for observation or analysis. Analytical instruments, comprising the spectrophotometer, electrophoresis apparatus and thermal cycler, generate a quantitative or molecular readout from a prepared sample. Support instruments, comprising the balance, pH meter and water bath, maintain the physical and chemical conditions on which the other three categories depend.

A student able to place an unfamiliar instrument within this framework rapidly infers its likely function, its probable failure modes and the precautions appropriate to its use, even before consulting its specific operating manual. This classification recurs implicitly throughout the book: each chapter from Part B onward introduces one or more instruments and states, at the outset, which of these four functional categories it occupies.

### **General Laboratory Rules**

- Wear a laboratory coat, closed footwear and appropriate eye protection at all times during practical work.
- Read the full procedure before starting; identify hazardous reagents and their first-aid measures in advance.
- Label every container, slide and reagent bottle immediately upon preparation.
- Never pipette any liquid by mouth.
- Report breakages, spills or injuries to the supervisor without delay.
- Dispose of biological, chemical and sharp waste in the designated containers, not in the general bin.
- Wash hands thoroughly before leaving the laboratory.

### **Quality Assurance and Reproducibility**

A technique carries scientific value only if it is reproducible, meaning that a second worker following the same written procedure on comparable material obtains a comparable result. Reproducibility depends on standardization at three levels: the reagent, prepared to a stated concentration and stored under stated conditions; the instrument, calibrated at a stated interval against a known reference; and the procedure itself, followed to a stated tolerance in timing and temperature.

Deviation at any one level introduces a source of error that cannot be separated, after the fact, from genuine biological variation. A competent practical record therefore states not only the result obtained but also the specific reagent batch, instrument and timing used to obtain it, so that an anomalous result can be traced to its likely origin rather than dismissed, or worse, accepted uncritically.


### **Case Illustration: Tracing a Data Error**

A student obtains an unexpectedly high protein estimate for a set of six tissue homogenates and, before reporting the value, checks the chain of technique behind it rather than accepting the number at face value. The calibration curve is found to be correctly linear, the spectrophotometer blank is correctly zeroed, but one reagent bottle used for the assay is discovered to be unlabelled with its preparation date. Because the working reagent for this assay

is known to degrade within a week of preparation, the unlabelled bottle cannot be confirmed as fresh, and the entire batch of six readings is repeated with a newly prepared, dated reagent rather than reported as obtained. The corrected values differ meaningfully from the first set. The episode illustrates directly why Table 1.1 treats labelling as a precaution rather than a formality: an unlabelled reagent converts an otherwise sound analytical procedure into an unverifiable one.

**Table 1.1 Common Laboratory Hazard Classes and Precautions**

| Hazard Class | Representative Example                   | Precaution                                                               |
|--------------|------------------------------------------|--------------------------------------------------------------------------|
| Chemical     | Concentrated acids; formalin vapour      | Use a fume hood; wear nitrile gloves and goggles                         |
| Biological   | Pathogenic cultures; fresh blood samples | Use personal protective equipment; autoclave waste before disposal       |
| Physical     | Broken glassware; heated equipment       | Handle with care; use heat-resistant gloves near ovens and autoclaves    |
| Electrical   | Frayed cords on instruments              | Inspect equipment before use; keep hands dry near sockets                |
| Fire         | Flammable solvents near an open flame    | Store solvents away from heat sources; keep an extinguisher within reach |



**Figure 1.1 General Workflow of a Zoological Investigation**

**QUICK RECAP**

- Technique is a reproducible procedure that converts biological material into verifiable evidence.
- Laboratory work follows a fixed sequence: preparation, execution, observation, recording.
- Personal protective equipment is mandatory against chemical, biological, physical, electrical and fire hazards.
- Every specimen, slide and reagent must be labelled at the moment of preparation.

**COMMON ERRORS TO AVOID**

- Beginning a procedure without reading it in full, leading to missed timing steps.
- Leaving slides, tubes or specimens unlabelled during a busy practical session.
- Treating chemical waste and general waste as interchangeable.

**Review Questions**

[UG] 1. Define zoological technique and state the four broad domains it covers.

[UG] 2. List five general laboratory safety rules applicable to a zoology practical class.

[PG] 3. Explain why incomplete recording of observations is considered a methodological failure rather than a clerical one.

[PG] 4. Classify the hazard types encountered in a histology laboratory and justify the precaution corresponding to each.

## CHAPTER 2. COLLECTION, PRESERVATION AND MUSEUM TECHNIQUES

*Learning focus: Field-heavy, strong UG core*

### Principle

Collection is the controlled removal of specimens from their natural habitat for scientific study. Preservation is the arrest of post-mortem decomposition so that morphological and, where relevant, molecular information remains available for later examination. The two steps are inseparable in practice: a specimen collected without an immediate plan for fixation is a specimen lost to autolysis and bacterial decay within hours.

### Representative Field Collection Equipment

- Aquatic fauna: plankton nets, dip nets, sieves, dredges, hand nets for shallow-water invertebrates.
- Terrestrial fauna: sweep nets and pitfall traps for arthropods, hand collection with forceps for slow-moving invertebrates, live traps for small mammals.
- Aerial fauna: aerial nets for flying insects, mist nets for birds and bats under appropriate permits.

### Killing and Narcotizing Agents

- Ethyl acetate vapour in killing jars produces rapid death in insects with minimal damage to setae and scales.
- Menthol crystals or carbon dioxide narcosis relax molluscs and other soft-bodied invertebrates before fixation, preventing muscular contraction that would distort the body.
- Anaesthetic overdose under institutional ethical clearance is required for vertebrate specimens.

### Permits and Ethical Collection

Collection of wild animal material in India is regulated under the Wildlife (Protection) Act and, for material of research or commercial value, under the Biological Diversity Act, both of which require prior permission from the relevant state forest department or the National Biodiversity Authority before a scheduled species is collected, transported or exported. A collector working



outside a protected area still requires landowner consent and, for an academic project, institutional ethical clearance describing the species, number of specimens and method of killing proposed.

Securing the appropriate permit before fieldwork begins is not a procedural formality: specimens collected without authorization cannot ordinarily be deposited in a recognized museum collection or cited in a peer-reviewed publication. The permit itself becomes part of the specimen's permanent scientific record alongside its locality label.

### **Fixation and Preservation**

Fixation denatures cellular protein and halts autolytic enzyme activity, stabilizing tissue architecture. Preservation, applied after fixation, maintains the specimen in that stabilized state over the long term. Fixative choice depends on the taxon, the intended downstream use and the duration of storage; Table 2.1 summarizes fixatives frequently used in zoological practice.

### **Wet Preservation versus Dry Preservation**

Wet preservation, in which the specimen is stored submerged in a preservative fluid, suits soft-bodied animals such as annelids, molluscs and amphibians. Dry preservation, comprising taxidermy, skeletal mounting and pinned insect collections, suits specimens with a rigid exoskeleton or those from which a durable mount can be prepared.

### **Outline Procedure for Preparing a Study Skin**

1. Record body measurements and locality data before any incision is made.
2. Make a mid-ventral incision; carefully separate the skin from the underlying musculature.
3. Remove adherent fat and tissue from the inner surface of the skin.
4. Treat the inner skin surface with a preservative powder such as borax.
5. Fill the skin with a suitable stuffing material, restoring natural body proportions.
6. Position the mount; allow it to dry fully; label with species identity, locality, date and collector.

### **Insect Pinning and Spreading**

Insects intended for a reference collection are pinned through the thorax using an entomological pin of appropriate gauge. Wings of Lepidoptera and Odonata are spread and dried on a setting board before the pin is transferred to a permanent storage drawer.

## Labelling, Cataloguing and Museum Management

A specimen without a locality label has severely reduced scientific value irrespective of its physical condition. Each specimen accessioned into a reference collection is assigned a unique catalogue number linked to a register recording species identity, collection locality, date, collector name and identification authority. Museum storage further requires control of temperature, humidity, light and pest infestation to prevent long-term deterioration of wet and dry material alike.

### Common Causes of Preservative Deterioration

A wet-preserved specimen deteriorates when the preservative fluid is allowed to evaporate below the level of the specimen, when the fluid is not renewed after it becomes diluted by residual tissue water, or when the storage container is exposed to strong light, which bleaches pigmentation and embrittles certain plastics over time. Routine museum maintenance therefore includes periodic inspection of fluid level and clarity, replacement of a cloudy or diluted preservative, and storage away from direct sunlight in a temperature-controlled room.


Dry-preserved material faces a different threat, principally insect pests such as dermestid beetles that consume skin, feather and fur. Regular inspection, an appropriate repellent, and quarantine of newly accessioned specimens before they join the main collection are standard preventive measures. Both preservation classes ultimately depend on the same principle introduced in Chapter 1: a specimen remains a scientific record only for as long as its physical integrity and its accompanying documentation are both maintained.

### Worked Example: Fixative Selection for a Field Survey

A survey team collecting frogs from a wetland intends to record external morphology, deposit voucher specimens in a reference museum, and additionally section the liver of a subset of individuals for histopathological screening. A single fixation protocol cannot serve all three purposes equally well: 10% formalin gives excellent long-term external preservation for the voucher series, but its slow penetration and moderate tissue hardening make it a suboptimal choice for the fine nuclear detail histopathology requires. The team therefore fixes the full voucher series in 10% formalin as planned, while the liver samples destined for histology are dissected out immediately upon collection and placed directly into Bouin's fluid, which penetrates the small tissue block rapidly and yields superior nuclear staining once processed as described in Chapter 4. Both fixatives are drawn from Table 2.1, chosen not by habit but by matching each fixative's known properties to the specific downstream use intended for that portion of the sample.

**Table 2.1 Common Fixatives Used in Zoological Preservation**

| Fixative                      | Composition                            | Typical Use                               | Approximate Duration                         |
|-------------------------------|----------------------------------------|-------------------------------------------|----------------------------------------------|
| Formalin (10%)                | Formaldehyde diluted in water          | General specimen preservation             | Several days, then transfer to storage fluid |
| Bouin's fluid                 | Picric acid, formaldehyde, acetic acid | Histological fixation of small specimens  | 12 to 24 hours                               |
| 70% Ethyl alcohol             | Ethanol diluted in water               | Long-term wet preservation                | Permanent, with periodic renewal             |
| FAA (Formalin-Acetic-Alcohol) | Formalin, acetic acid, ethanol         | Preservation of soft-bodied invertebrates | 24 to 48 hours                               |
| Carnoy's fluid                | Ethanol, chloroform, acetic acid       | Chromosome and nuclear fixation           | 1 to 3 hours                                 |

**Figure 2.1 Preservation and Cataloguing Pathway of a Field-collected Specimen****QUICK RECAP**

- Fixation halts decomposition; preservation maintains the fixed state over time.
- Fixative choice depends on taxon, downstream use and storage duration.
- Wet preservation suits soft-bodied animals; dry preservation suits animals with rigid support structures.
- A specimen without accurate locality and date data has diminished scientific value.

**COMMON ERRORS TO AVOID**

- Delaying fixation after collection, allowing autolysis to begin.
- Using an excessive formalin concentration, which hardens tissue and hinders later histological processing.
- Omitting locality or date information from the specimen label.

**Review Questions**

[UG] 1. Distinguish between fixation and preservation, giving one example of each.

[UG] 2. Describe the outline procedure for preparing a study skin of a small vertebrate.

[PG] 3. Justify the selection of Bouin's fluid over formalin for a specimen intended for histological sectioning.

[PG] 4. Discuss the minimum data required on a museum specimen label and its bearing on later taxonomic revision.

# CHAPTER 3. MICROSCOPY: PRINCIPLES AND APPLICATIONS

*Learning focus: UG basics; PG extends to SEM, TEM and confocal imaging*

## Principle of Light Microscopy

A compound light microscope forms a magnified image of a specimen by passing light sequentially through a condenser, the specimen, an objective lens and an ocular lens. Total magnification is the product of the objective and ocular magnifications. Resolving power, the ability to distinguish two close points as separate, depends on the wavelength of light used and the numerical aperture of the objective rather than on magnification alone.

## The Compound and Stereo Microscope

The compound microscope is used for thin, translucent preparations such as stained sections and smears. The stereo, or dissecting, microscope provides a lower-magnification three-dimensional view suited to gross dissection and sorting of whole specimens, fitted with a wider working distance for manipulation under the lens.

## Illumination and Köhler Alignment

Image quality in light microscopy depends as much on correct illumination as on the lens system itself. Köhler illumination, the standard alignment used in research microscopes, focuses the light source at the plane of the condenser aperture rather than at the specimen plane, producing even, glare-free illumination across the field of view regardless of the lamp filament's own structure. Correct condenser height and aperture diaphragm setting, adjusted for each objective in turn, control both illumination evenness and the effective numerical aperture reaching the specimen; closing the aperture diaphragm too far increases apparent contrast but reduces true resolving power, a trade-off students frequently mismanage when a preparation appears insufficiently sharp.

## Routine Care of a Compound Microscope

- Carry the instrument with both hands, one supporting the base.
- Clean lenses only with lens tissue, never with a laboratory cloth.
- Begin focusing on the lowest objective power before switching to higher magnification.
- Cover the microscope after use; store it in a dust-free cabinet.

## **Phase-contrast and Fluorescence Microscopy**

Phase-contrast microscopy converts small differences in refractive index within an unstained, living specimen into visible differences in brightness, allowing observation of living cells without fixation or staining. Fluorescence microscopy illuminates a specimen labelled with a fluorochrome using light of a specific excitation wavelength, then detects the longer-wavelength light re-emitted by the fluorochrome, giving high-contrast localization of the labelled molecule against a dark background.

## **Confocal Laser Scanning Microscopy**

Confocal microscopy uses a focused laser beam and a pinhole aperture placed before the detector to exclude out-of-focus light, so that only light from a single thin optical plane contributes to the image. Successive optical sections through a fluorescently labelled specimen can be reconstructed into a three-dimensional image, a property valuable for thick tissue and whole-mount preparations.

## **Electron Microscopy**

Electron microscopes use a beam of accelerated electrons in place of light, achieving resolution several orders of magnitude finer than any light-based instrument, because the wavelength of an electron beam is far shorter than that of visible light.

Scanning electron microscopy (SEM) directs a focused electron beam across the surface of a specimen and detects electrons scattered from that surface, producing a detailed three-dimensional image of surface topography. Specimens require fixation, dehydration, critical-point drying and coating with a thin conductive layer, typically gold or platinum, before examination.

Transmission electron microscopy (TEM) passes an electron beam through an ultrathin section, typically 60 to 90 nanometres thick, forming an image from the electrons transmitted through the specimen. TEM resolves internal cellular ultrastructure, including organelle membranes, beyond the reach of light microscopy.

## **Selecting a Microscopy Technique for a Research Question**

The choice among the techniques compared in Table 3.1 follows from the biological question asked rather than from equipment availability alone. A question about living cell behaviour, such as ciliary movement or phagocytosis, requires phase-contrast or, where a specific protein must be tracked, fluorescence microscopy on a live, unfixed preparation. A question about the three-dimensional distribution of a labelled structure within a thick tissue block is best addressed by confocal microscopy, which alone permits optical sectioning without physical cutting.

A question about external surface architecture, such as insect cuticular ornamentation, calls for scanning electron microscopy, whereas a question about internal organelle structure, such as mitochondrial cristae morphology, requires transmission electron microscopy. Selecting a lower-resolution, lower-cost technique adequate to the question asked is good practice; selecting a technique inadequate to the question is a common and avoidable design error in student projects.

### Worked Example: Calculating Total Magnification


A student examines a stained blood smear using a 40x objective and a 10x ocular lens. Total magnification is the product of the two values: 40 multiplied by 10 gives 400x, meaning a structure 400 times its true size appears on the retina or camera sensor. Switching to a 100x oil-immersion objective with the same 10x ocular raises total magnification to 1000x, the setting conventionally used to examine red blood cell morphology and to search for intracellular parasites such as *Plasmodium*. Magnification alone, however, does not guarantee that finer detail becomes visible: if the objective's numerical aperture is insufficient for the structure in question, increasing magnification further only enlarges an already blurred image, described among microscopists as "empty magnification," rather than resolving any additional true detail, which is why Table 3.1 reports resolution as a property distinct from, and ultimately more informative than, magnifying power alone.

**Table 3.1 Comparison of Microscopy Techniques**

| Technique                          | Approximate Resolution | Sample Requirement                    | Typical Use                          |
|------------------------------------|------------------------|---------------------------------------|--------------------------------------|
| Light (compound) microscopy        | 0.2 micrometre         | Thin stained section or smear         | Routine histology, cytology          |
| Phase-contrast microscopy          | 0.2 micrometre         | Unstained living cells                | Observation of living cell structure |
| Fluorescence microscopy            | 0.2 micrometre         | Fluorochrome-labelled specimen        | Localization of specific molecules   |
| Confocal laser scanning microscopy | 0.1 to 0.2 micrometre  | Fluorescently labelled thick specimen | Three-dimensional cellular imaging   |
| Scanning electron microscopy       | 1 to 10 nanometre      | Metal-coated dry specimen             | Surface ultrastructure               |



| Technique                        | Approximate Resolution | Sample Requirement      | Typical Use             |
|----------------------------------|------------------------|-------------------------|-------------------------|
| Transmission electron microscopy | 0.1 to 0.2 nanometre   | Ultrathin resin section | Internal ultrastructure |



**Figure 3.1 Path of Light in a Compound Microscope**

### QUICK RECAP

- Magnification is the product of objective and ocular power; resolution depends on wavelength and numerical aperture.
- Phase-contrast microscopy permits observation of living, unstained cells.
- Confocal microscopy excludes out-of-focus light, permitting optical sectioning of thick specimens.
- SEM images surface topography; TEM images internal ultrastructure, both at resolutions beyond light microscopy.

### COMMON ERRORS TO AVOID

- Focusing on high power before locating the specimen on low power, risking slide damage.
- Cleaning objective lenses with an unsuitable cloth, scratching the lens coating.
- Confusing magnification with resolving power when comparing instruments.

### Review Questions

**[UG]** 1. State the formula for total magnification in a compound microscope and define resolving power.

**[UG]** 2. Differentiate between a compound microscope and a stereo microscope in terms of use.

**[PG]** 3. Explain the role of the pinhole aperture in confocal laser scanning microscopy.

**[PG]** 4. Compare the sample preparation requirements of SEM and TEM, relating each to the type of structural information obtained.

# Chapter 4. Microtomy and Histological Techniques

## PART B · HISTOLOGICAL AND CYTOLOGICAL TECHNIQUES

*From tissue to slide: processing, staining and chromosome analysis*

**Learning focus:** Core UG and PG practical skill

### Principle

Histological technique converts a piece of solid tissue into a thin, stained section suitable for microscopic examination. The pathway comprises fixation, dehydration, clearing, infiltration, embedding, sectioning, staining and mounting, each step preparing the tissue for the one that follows. Failure at any single step, most commonly incomplete fixation or incomplete dehydration, compromises every subsequent step and cannot be corrected later.

### Fixation

Fixation cross-links or precipitates cellular protein, arresting autolysis and preserving structural relationships between cell components. Common fixatives include 10 percent neutral buffered formalin for general histology, Bouin's fluid for delicate tissue, and glutaraldehyde for material destined for electron microscopy. Fixative volume should exceed tissue volume by a factor of fifteen to twenty to ensure adequate penetration.

### Choice of Fixative and Common Artefacts

Fixative selection balances speed of penetration against the degree of chemical alteration introduced into the tissue. Formalin penetrates slowly but preserves antigenicity well, favouring its use where a section may later require immunohistochemical labelling. Bouin's fluid penetrates rapidly and gives excellent nuclear detail in small specimens, but the picric acid it contains can interfere with certain downstream molecular applications unless thoroughly washed out.

Common fixation artefacts include shrinkage from prolonged exposure to a strongly dehydrating fixative, and a hardened, difficult-to-section block from a fixative left to act for far longer than necessary. Recognizing an artefact as a technical error, rather than a genuine tissue feature, is a skill developed only through repeated comparison of a poorly and a well-processed section of the same tissue type.

## Tissue Processing Sequence

1. Dehydration: the tissue passes through an ascending series of ethanol grades, typically 50, 70, 90 and 100 percent, removing tissue water.
2. Clearing: a clearing agent such as xylene replaces the dehydrating alcohol and renders the tissue translucent, at the same time being miscible with the embedding medium.
3. Infiltration: molten paraffin wax replaces the clearing agent within the tissue, providing internal support for sectioning.
4. Embedding: the infiltrated tissue is oriented and cast into a wax block, which is allowed to solidify completely before sectioning.

## Sectioning

A rotary microtome cuts paraffin sections between four and six micrometres in thickness for routine work. A cryostat, which sections frozen unfixed or lightly fixed tissue at sub-zero temperature, is used where speed is required or where a reagent under study would be lost during paraffin processing, such as certain enzymes or lipids. Sections are floated on a warm water bath to remove cutting compression, then mounted on a glass slide.

## Staining and Mounting

Haematoxylin and eosin (H&E) is the routine stain of histology: haematoxylin, a basic dye, colours acidic structures such as the nucleus blue to purple, while eosin, an acidic dye, colours basic cytoplasmic and extracellular structures pink. Special stains extend this baseline where a specific tissue component is of interest, as summarised for histochemical methods in Chapter 5. Following staining, the section is dehydrated, cleared and mounted permanently under a coverslip using a resinous mounting medium.

## Interpreting a Routine Section

A well-prepared H&E section should show intact tissue architecture free of folds, tears or air bubbles beneath the coverslip, with the nucleus sharply differentiated in blue-purple against pink cytoplasmic and stromal detail. Uneven staining across a section commonly indicates incomplete deparaffinization or an exhausted staining reagent rather than a true biological difference, and the section should be repeated before any structural conclusion is drawn from it. Systematic scanning of the section at low power before selecting a field for high-power examination avoids the common beginner error of describing an unrepresentative, atypical region of the tissue as though it were characteristic of the whole.


## Worked Example: Diagnosing a Processing Fault

A batch of liver sections examined after routine H&E staining shows widespread cell shrinkage, with clear, empty-looking spaces separating the cells from one another across the entire slide.

Because the fault appears uniformly across every section in the batch rather than in a single section, a fault in an individual step of processing, rather than a peculiarity of one animal, is the more likely explanation. Reviewing the processing log for the batch reveals that the ninety and hundred percent ethanol stations had not been refreshed for several weeks and had become diluted through carryover from earlier batches, so that dehydration was effectively incomplete despite the tissue having passed through the full nominal schedule in Table 4.1. Replacing the ethanol series and reprocessing a fresh batch from the same fixed tissue block resolves the artefact, confirming the diagnosis and illustrating why a processing schedule specifies not only duration but also the maintained quality of each reagent station.

**Table 4.1 Representative Tissue Processing Schedule**

| Step         | Reagent                              | Approximate Duration | Purpose                                         |
|--------------|--------------------------------------|----------------------|-------------------------------------------------|
| Fixation     | 10% neutral buffered formalin        | 12 to 24 hours       | Preservation of tissue structure                |
| Dehydration  | Ascending ethanol series, 50 to 100% | 30 minutes per grade | Removal of tissue water                         |
| Clearing     | Xylene                               | 30 to 45 minutes     | Removal of dehydrating agent                    |
| Infiltration | Molten paraffin wax                  | 1 to 2 hours         | Support for sectioning                          |
| Embedding    | Paraffin block                       | Until solidified     | Formation of the tissue block                   |
| Sectioning   | Rotary microtome                     | Variable             | Production of thin sections, 4 to 6 micrometres |



**Figure 4.1 Histological Technique Pipeline**

**QUICK RECAP**

- The histological pathway proceeds through fixation, dehydration, clearing, infiltration, embedding, sectioning, staining and mounting.
- Fixative volume should exceed tissue volume fifteen to twenty times to ensure adequate penetration.
- Routine paraffin sections are cut at four to six micrometres; a cryostat sections frozen tissue for speed or heat-labile components.
- H&E staining differentiates nucleus (haematoxylin, basic dye) from cytoplasm (eosin, acidic dye).

**COMMON ERRORS TO AVOID**

- Under-fixing a large tissue block, leaving the core poorly preserved.
- Incomplete dehydration, leaving residual water that produces poor sectioning quality.
- Sectioning before the wax block has fully hardened, producing torn or compressed ribbons.

**Review Questions**

[UG] 1. List the sequence of steps in routine paraffin histological technique.

[UG] 2. State the colour reactions of haematoxylin and eosin and the structures each stains.

[PG] 3. Explain why a cryostat section is preferred over a paraffin section for the demonstration of certain enzymes.

[PG] 4. Discuss the consequences of inadequate fixative volume relative to tissue volume.

# Chapter 5. Histochemistry and Immunohistochemistry

*Learning focus: PG-weighted; UG overview*

## Principle of Histochemistry

Histochemistry localizes a specific chemical constituent within a tissue section through a reaction that yields a visible, usually coloured, end product at the site of the target molecule. Unlike routine staining, which distinguishes tissue components largely by dye affinity, a histochemical reaction is chosen for its chemical specificity for carbohydrate, lipid, protein, nucleic acid or enzyme activity.

## Representative Histochemical Reactions

- The periodic acid-Schiff (PAS) reaction oxidizes vicinal diols in neutral carbohydrates to aldehydes, which react with Schiff reagent to give a magenta product; it demonstrates glycogen and mucosubstances.
- Sudan Black B dissolves selectively in tissue lipid, producing a black colouration used to demonstrate fat deposits.
- The Feulgen reaction hydrolyses DNA to expose aldehyde groups that react with Schiff reagent, giving a purple-red, DNA-specific colouration used in quantitative cytophotometry.
- Enzyme histochemistry captures the product of an enzyme-substrate reaction as an insoluble coloured precipitate at the exact site of enzyme activity, for example alkaline phosphatase localization.

## Choosing a Histochemical Method

The histochemical methods listed in Table 5.1 are chosen according to the tissue component under investigation, since no single stain is informative for every biomolecule class. A study of glycogen storage in an invertebrate hepatopancreas, for example, is appropriately investigated with the periodic acid-Schiff reaction rather than a general connective-tissue stain, while a study of lipid mobilization during starvation is appropriately investigated with Sudan Black B on frozen, unprocessed tissue, because the organic solvents used in routine paraffin processing would otherwise dissolve the very lipid the study seeks to demonstrate.

## **Principle of Immunohistochemistry**

Immunohistochemistry (IHC) exploits the specific binding of an antibody to its target antigen to localize a single protein within a tissue section. In the indirect method, an unlabelled primary antibody binds the target antigen, and a labelled secondary antibody directed against the primary antibody then binds in turn; the label, typically an enzyme such as horseradish peroxidase, converts a colourless substrate to a coloured, insoluble product at the site of binding. Because several secondary antibody molecules can bind each primary antibody, the indirect method amplifies the signal relative to a directly labelled primary antibody.

## **Antigen Retrieval and Fluorescence In-Situ Hybridization**

Formalin fixation cross-links protein and can mask antigenic sites; antigen retrieval, achieved by heating the section in a buffer or by limited proteolytic digestion, restores antibody accessibility before the IHC procedure. Fluorescence in-situ hybridization (FISH) applies a comparable principle to nucleic acid: a fluorochrome-labelled DNA probe hybridizes to a complementary sequence within the tissue or chromosome preparation, permitting direct visualization of a specific gene locus.

## **Controls in Histochemistry and Immunohistochemistry**


Interpretation of any histochemical or immunohistochemical result depends absolutely on an appropriate control run alongside the test section. A positive control, a section known to contain the target constituent, confirms that the reagents and procedure performed as expected; a negative control, in which the primary antibody or key reagent is omitted, confirms that the observed colouration reflects the intended reaction rather than non-specific background binding. A result reported without a matched control accompanying it cannot be regarded as scientifically interpretable, however striking the colouration obtained.

## **Case Illustration: Confirming Glycogen with a Diastase Control**

A section stained by the periodic acid-Schiff method shows strong magenta granules within hepatocyte cytoplasm, consistent with glycogen storage, but PAS alone cannot distinguish glycogen from other neutral carbohydrate material that also reacts positively. To confirm the identification, a duplicate, adjacent section is first digested with the enzyme diastase, which specifically hydrolyses glycogen, before PAS staining is applied. Loss of the magenta reaction in the diastase-treated section, contrasted against its retention in the untreated section, confirms that glycogen specifically, rather than another PAS-positive substance, accounted for the original staining. This paired diastase-digestion control is the standard method by which a PAS result is upgraded from a suggestive observation to a specific, defensible identification.

**Table 5.1 Histochemical Stains and Target Biomolecules**

| Stain / Method                      | Target Biomolecule              | Typical Result                        |
|-------------------------------------|---------------------------------|---------------------------------------|
| Periodic acid-Schiff (PAS)          | Neutral carbohydrates, glycogen | Magenta colouration                   |
| Sudan Black B                       | Lipids                          | Black colouration                     |
| Feulgen reaction                    | DNA                             | Purple-red colouration                |
| Mercury bromophenol blue            | Proteins                        | Blue colouration                      |
| Alkaline phosphatase histochemistry | Enzyme activity site            | Coloured precipitate at reaction site |

**Figure 5.1 Principle of Indirect Immunohistochemistry****QUICK RECAP**

- Histochemistry localizes a chemical constituent through a reaction specific to that constituent.
- PAS demonstrates carbohydrate; Sudan Black B demonstrates lipid; the Feulgen reaction is DNA-specific.
- Indirect immunohistochemistry uses a labelled secondary antibody to amplify signal from an unlabelled primary antibody.
- Antigen retrieval restores antibody accessibility lost during formalin fixation.



**COMMON ERRORS TO AVOID**

- Applying IHC without an antigen retrieval step, giving weak or absent staining.
- Omitting a negative control, making a background reaction indistinguishable from true positive labelling.
- Interpreting a histochemical colour reaction without a matched, appropriately stained control section.

**Review Questions**

[UG] 1. Explain the chemical basis of the periodic acid-Schiff reaction.

[UG] 2. State the purpose of the secondary antibody in indirect immunohistochemistry.

[PG] 3. Discuss the need for antigen retrieval in formalin-fixed, paraffin-embedded sections.

[PG] 4. Compare the principle of immunohistochemistry with that of fluorescence in-situ hybridization.

# Chapter 6. Chromosome Preparation and Karyotyping

*Learning focus: PG core; UG introductory*

## Principle

Chromosome preparation captures dividing cells at metaphase, the stage at which chromosomes are maximally condensed and individually distinguishable, then spreads them on a slide for microscopic analysis. The resulting karyotype, an ordered display of an organism's chromosome complement by size and morphology, is basic to cytotaxonomy, cytogenetic diagnosis and evolutionary comparison between related species.

## Steps in Metaphase Chromosome Preparation

1. Obtain actively dividing cells from bone marrow, peripheral blood culture, or a cultured cell line.
2. Arrest cells at metaphase using colchicine or a related spindle-inhibiting agent.
3. Treat cells with a hypotonic solution, causing them to swell and disperse the chromosomes.
4. Fix cells in a mixture of methanol and glacial acetic acid, typically in a three-to-one ratio.
5. Drop the fixed cell suspension onto a clean, chilled slide and allow it to air-dry, spreading the chromosomes in a single plane.
6. Stain the preparation and examine it under oil-immersion light microscopy.

## Sources of Dividing Cells and Culture Considerations

Bone marrow provides an actively dividing cell population directly, without the need for prior culture, and is favoured where a rapid preparation is required from a small mammal. Peripheral blood requires a short-term culture, typically 48 to 72 hours in a medium containing a mitogen such as phytohaemagglutinin to stimulate lymphocyte division, before the same colchicine-hypotonic-fixation sequence is applied. A cultured cell line offers the most reliable and repeatable source of metaphase plates, since its growth conditions and division rate are already standardized, but is unavailable for species not yet established in culture, a substantial limitation for comparative cytogenetic surveys of wild, non-model taxa.

## Chromosome Banding

Uniform staining, as with Giemsa alone, identifies chromosome number and gross morphology but cannot distinguish chromosomes of similar size. Banding techniques introduce a longitudinal pattern of light and dark bands unique to each chromosome pair, permitting unambiguous identification and detection of structural rearrangement.

## Karyotype Analysis and Polytene Chromosomes

A karyotype is constructed by photographing or digitally capturing a well-spread metaphase plate, then arranging the chromosomes in homologous pairs by decreasing size to produce an idiogram. Polytene chromosomes, found in certain insect tissues such as the dipteran salivary gland, arise through repeated DNA replication without cell division and display a characteristic banding pattern directly visible without artificial banding treatment, making them useful for fine-scale mapping of chromosomal rearrangement.

## Applications in Comparative Cytogenetics

Karyotype data support taxonomy directly where closely related species cannot be reliably separated by external morphology alone but differ in chromosome number or banding pattern, a situation common among rodents, fish and several insect groups. Comparative karyotyping across a genus can additionally reveal the chromosomal rearrangements, such as fusion, fission or inversion, that accompanied speciation, providing evolutionary evidence independent of, and often complementary to, the molecular sequence data described in Part D.


## Worked Example: Constructing an Idiogram

A well-spread metaphase plate from a rodent bone marrow preparation is photographed, and the length of each of its forty chromosomes is measured from the digital image along with the position of each centromere. Chromosomes are then paired by matching each measured length and centromere position against its closest counterpart, giving twenty homologous pairs, which are numbered from largest to smallest and arranged in a standard row for presentation as an idiogram. A pair whose centromere lies at the exact midpoint is classified as metacentric, one with the centromere distinctly off-centre as submetacentric, and one with the centromere near one end as acrocentric, a classification that itself frequently assists in tracking which pair corresponds to which chromosome across successive preparations of the same species.

**Table 6.1 Chromosome Banding Techniques Compared**

| Technique | Principle                                          | Primary Application                           |
|-----------|----------------------------------------------------|-----------------------------------------------|
| C-banding | Selective staining of constitutive heterochromatin | Identification of centromeric heterochromatin |

| Technique    | Principle                                      | Primary Application                         |
|--------------|------------------------------------------------|---------------------------------------------|
| G-banding    | Trypsin treatment followed by Giemsa staining  | Chromosome identification and karyotyping   |
| Q-banding    | Quinacrine fluorescent staining                | Fluorescent chromosome identification       |
| NOR staining | Silver staining of nucleolar organizer regions | Localization of ribosomal RNA gene clusters |



**Figure 6.1 Steps in Metaphase Chromosome Preparation**

### QUICK RECAP

- Metaphase is the chromosome stage used for routine chromosome preparation because condensation is maximal.
- Colchicine arrests mitosis; hypotonic treatment disperses chromosomes for clean spreading.
- Banding techniques permit identification of individual chromosome pairs beyond size and shape alone.
- Polytene chromosomes display a natural banding pattern useful for fine chromosomal mapping.

### COMMON ERRORS TO AVOID

- Insufficient hypotonic treatment, leaving chromosomes clumped and difficult to score.
- Excessive colchicine exposure, causing chromosome over-contraction.
- Dropping the fixed suspension onto a warm or greasy slide, giving poor spreading.

### Review Questions

[UG] 1. Define karyotype and state the metaphase stage's relevance to its preparation.

- [UG]** 2. List the sequential steps of metaphase chromosome preparation.
- [PG]** 3. Differentiate C-banding from G-banding in principle and application.
- [PG]** 4. Explain why polytene chromosomes are useful in fine-scale chromosomal mapping.

# Chapter 7. Centrifugation Techniques

## PART C · BIOPHYSICAL AND BIOCHEMICAL TECHNIQUES

*Instrumentation for separating, quantifying and analysing biomolecules*

**Learning focus:** PG-weighted; UG overview

### Principle

Centrifugation separates particles suspended in a liquid according to their size, shape and density by subjecting the suspension to a centrifugal field generated by rapid rotation. The effective force is expressed as relative centrifugal force (RCF), a multiple of standard gravity that depends on rotational speed and the radius of rotation; larger or denser particles sediment at lower speed than smaller or less dense ones.

### Sedimentation Behaviour

A particle in a centrifugal field sediments at a rate governed by its mass, its shape and the difference between its own density and that of the surrounding medium; a dense, compact particle such as a nucleus sediments far more readily than a diffuse, low-density structure such as a microsomal vesicle. This differential behaviour, rather than any active separation mechanism, is what allows a mixed cellular homogenate to be resolved into its component fractions purely through a sequence of centrifugal steps of increasing severity.

### Types of Centrifuges and Rotors

- Low-speed clinical centrifuges sediment whole cells and coarse debris.
- High-speed centrifuges sediment organelles and coarse precipitates.
- Ultracentrifuges, operating at the highest speeds, separate macromolecules such as ribosomes and viral particles, and support density-gradient work.
- Fixed-angle rotors hold tubes at a constant angle, giving rapid pelleting; swinging-bucket rotors reorient during the run, giving cleaner gradient separation.

### Differential and Density-Gradient Centrifugation

Differential centrifugation subjects a homogenate to a series of successively increasing speeds; at each step, particles heavier than the prevailing threshold sediment as a pellet while lighter particles remain in the supernatant, which is then carried forward to the next, faster spin. Density-gradient centrifugation instead layers the sample over a preformed gradient of increasing density, typically sucrose or caesium chloride, and separates particles either by the

depth at which their own density equals that of the surrounding medium, or by their sedimentation rate through the gradient.

### Precautions in Centrifuge Operation


Tubes loaded into a rotor must be balanced in matched, opposite pairs of equal mass, since an unbalanced rotor spinning at high speed generates a vibration capable of damaging the drive shaft or, in a severe case, causing rotor failure. Ultracentrifuge rotors additionally carry a defined maximum speed rating that must never be exceeded, and are retired from service after a stated number of runs regardless of apparent condition, because metal fatigue accumulates invisibly with repeated stress at high rotational force.

### Worked Example: Calculating Relative Centrifugal Force

Relative centrifugal force is calculated as 1.118 multiplied by 10 to the power of minus five, multiplied by the rotor radius in centimetres, multiplied by the square of the rotation speed in revolutions per minute. For a rotor of radius 10 centimetres spinning at 3,000 rpm, RCF equals 1.118 times 10 to the power of minus five, times 10, times 3,000 squared, which works out to approximately 1,006 times gravity, sufficient to pellet whole cells and nuclei but well below the force needed to sediment ribosomes or viral particles. Doubling the rotation speed to 6,000 rpm at the same radius raises RCF roughly fourfold rather than twofold, since force scales with the square of speed, a relationship that explains why a modest increase in rpm produces a disproportionately large increase in sedimenting force.

**Table 7.1 Common Centrifuge Types and Speed Ranges**

| Type                            | Approximate Speed Range    | Typical Application                                     |
|---------------------------------|----------------------------|---------------------------------------------------------|
| Low-speed (clinical) centrifuge | Up to 6,000 rpm            | Sedimentation of cells, coarse debris                   |
| High-speed centrifuge           | 6,000 to 25,000 rpm        | Sedimentation of organelles, precipitates               |
| Ultracentrifuge                 | 25,000 to over 100,000 rpm | Separation of macromolecules, subcellular fractionation |



**Figure 7.1 Differential Centrifugation Scheme for Subcellular Fractionation**

### QUICK RECAP

- Relative centrifugal force depends on rotational speed and radius, not on speed alone.
- Differential centrifugation separates particles stepwise by size and density using successive speed increases.
- Density-gradient centrifugation resolves particles along a preformed density gradient.
- Fixed-angle rotors favour speed of pelleting; swinging-bucket rotors favour gradient resolution.

### COMMON ERRORS TO AVOID

- Loading unbalanced tubes, causing rotor vibration or failure.
- Selecting a differential protocol when a density-gradient separation is required for the resolution needed.
- Disturbing the pellet while decanting the supernatant.

### Review Questions

[UG] 1. Define relative centrifugal force and the factors on which it depends.

[UG] 2. Distinguish a fixed-angle rotor from a swinging-bucket rotor.

[PG] 3. Outline the sequence of a differential centrifugation protocol used for subcellular fractionation.

[PG] 4. Compare density-gradient centrifugation with differential centrifugation in terms of resolving power.



# Chapter 8. Spectrophotometry and Colorimetry

*Learning focus: Common to UG and PG laboratories*

## Principle: The Beer-Lambert Law

Spectrophotometry measures the amount of light of a defined wavelength absorbed by a solution, relating that absorbance quantitatively to the concentration of the absorbing substance through the Beer-Lambert law,  $A$  equals  $\epsilon$  multiplied by  $c$  multiplied by  $l$ , where  $\epsilon$  is the molar extinction coefficient,  $c$  is concentration and  $l$  is the path length of light through the sample. Because absorbance is directly proportional to concentration at a fixed path length, a calibration curve prepared from known standards allows the concentration of an unknown sample to be read from its measured absorbance.

## The Calibration Curve

Quantification by spectrophotometry rests on a calibration curve prepared from a dilution series of a known standard, with absorbance plotted against concentration; because the Beer-Lambert relationship is linear only over a limited concentration range, an unknown sample giving an absorbance value outside the linear portion of the curve must be diluted and re-measured rather than extrapolated, since extrapolation beyond the calibrated range systematically overestimates or underestimates the true concentration.

## Instrumentation

A UV-Vis spectrophotometer directs light from a source through a monochromator, which selects a narrow band of wavelength, through the sample cuvette, to a photodetector that converts transmitted light intensity into an electrical signal. A single-beam instrument measures the blank and the sample sequentially through one optical path; a double-beam instrument splits the light into a reference path and a sample path measured simultaneously, giving greater stability against source fluctuation.

## Representative Colorimetric Applications in Zoology

- Protein estimation by the Biuret or Bradford method, read at a fixed visible wavelength against a bovine serum albumin standard curve.
- Nucleic acid quantification by absorbance at 260 nanometres, with the ratio of absorbance at 260 to 280 nanometres used to assess protein contamination.

- Enzyme assays in which the rate of formation of a coloured product is monitored over time to determine enzyme activity.

### Flame Photometry

Flame photometry, a related technique, atomizes a sample in a flame and measures the intensity of characteristic light emitted by specific metal ions, most commonly sodium and potassium, providing rapid estimation of these electrolytes in biological fluids.

### Sources of Error


Stray light, an unclean cuvette, an air bubble in the light path and an incorrectly matched reagent blank are the principal sources of error in routine spectrophotometric work. Because a small error in absorbance translates directly into a proportional error in the reported concentration, technique that appears merely careless, such as handling a cuvette by its optical faces, has a direct and quantifiable effect on data quality rather than being a purely cosmetic lapse.

### Worked Example: Reading Concentration from a Calibration Curve

A calibration series of bovine serum albumin standards at 0, 20, 40, 60, 80 and 100 micrograms per millilitre gives absorbance readings of 0.00, 0.21, 0.42, 0.63, 0.84 and 1.05 respectively at 595 nanometres, a straight-line relationship confirming the assay is behaving as the Beer-Lambert law predicts across this range. An unknown tissue extract gives an absorbance reading of 0.53 under identical conditions; reading this value against the calibration line, or equivalently dividing 0.53 by the slope of approximately 0.0105 absorbance units per microgram per millilitre, gives an estimated concentration of approximately 50 micrograms per millilitre. Had the unknown instead given a reading of 1.40, outside the calibrated range entirely, the correct response would be to dilute the extract and re-assay it, not to extrapolate the line beyond the concentrations actually tested.

**Table 8.1 Beer-Lambert Law: Variables and Units**

| Symbol  | Quantity                            | Unit                          |
|---------|-------------------------------------|-------------------------------|
| A       | Absorbance                          | Dimensionless                 |
| epsilon | Molar extinction coefficient        | Litre per mole per centimetre |
| c       | Concentration of solution           | Mole per litre                |
| l       | Path length of light through sample | Centimetre                    |



**Figure 8.1 Optical Path in a Single-beam UV-Vis Spectrophotometer**

### QUICK RECAP

- Absorbance is proportional to concentration at constant path length, per the Beer-Lambert law.
- A double-beam spectrophotometer corrects for source fluctuation by simultaneous reference measurement.
- The 260 to 280 nanometre absorbance ratio indicates protein contamination of a nucleic acid sample.
- Flame photometry estimates sodium and potassium through characteristic flame emission.

### COMMON ERRORS TO AVOID

- Failing to zero the instrument against an appropriate blank before sample measurement.
- Using a fingerprint-smudged or scratched cuvette, introducing stray light error.
- Reading absorbance outside the linear range of the calibration curve.

### Review Questions

[UG] 1. State the Beer-Lambert law and define each of its terms.

[UG] 2. Differentiate a single-beam from a double-beam spectrophotometer.

[PG] 3. Explain the significance of the 260 to 280 nanometre absorbance ratio in nucleic acid quality assessment.

[PG] 4. Describe the principle of flame photometry and its typical biological application.

## Chapter 9. Chromatographic Techniques

*Learning focus: PG core; UG introductory*

### Principle

Chromatography separates the components of a mixture by their differential distribution between a stationary phase and a moving mobile phase. Components with greater affinity for the stationary phase migrate more slowly than those with greater affinity for the mobile phase, so that a mixture applied at a single point resolves into spatially or temporally separated bands as the mobile phase advances.

### Paper and Thin-layer Chromatography

In paper chromatography, cellulose paper serves as the stationary phase and a liquid solvent as the mobile phase; the distance travelled by a compound relative to the solvent front, expressed as the retardation factor or  $R_f$  value, is used for tentative identification. Thin-layer chromatography (TLC) replaces paper with a thin layer of silica gel or alumina coated on a rigid support, offering faster separation and sharper resolution than paper chromatography for the qualitative screening of amino acids, lipids and pigments.

### Interpreting an $R_f$ Value

The  $R_f$  value of a compound is characteristic for a given stationary phase, mobile phase and temperature, but is not an absolute physical constant and cannot be compared reliably between runs performed under different conditions. Confident identification of an unknown compound therefore requires the unknown to be run alongside an authentic reference standard on the same plate under the same conditions, so that identity is judged by co-migration to an identical  $R_f$  rather than by comparison with a value quoted from a different laboratory or a different day.

## Column, Gas and High-performance Liquid Chromatography

Column chromatography packs an adsorbent into a vertical column and is used at preparative scale to isolate a purified fraction from a mixture. Gas chromatography (GC) vaporizes the sample and carries it through a coated capillary column by an inert carrier gas, resolving volatile compounds such as fatty acid methyl esters or pesticide residues by their differential retention within the column. High-performance liquid chromatography (HPLC) forces a liquid mobile phase through a densely packed column under high pressure, achieving high-resolution separation and quantification of non-volatile compounds such as proteins, hormones and pigments.

## Applications in Zoological Research


Chromatographic technique supports zoological research across pigment biology, where paper and thin-layer chromatography separate the carotenoid and pterin pigments responsible for animal colouration; toxicology, where gas chromatography detects and quantifies pesticide residue in tissue as introduced in Chapter 16; and endocrinology, where HPLC quantifies circulating hormone concentration from a small blood or tissue sample with a sensitivity that a colorimetric assay alone cannot match.

## Worked Example: Calculating an R<sub>f</sub> Value

A pigment spot applied near the base of a thin-layer plate migrates 6.4 centimetres from the origin while the solvent front migrates 8.0 centimetres before the plate is removed from the developing chamber. The R<sub>f</sub> value is the ratio of these two distances, 6.4 divided by 8.0, giving 0.80. A second, co-run reference standard of known identity also migrates to an R<sub>f</sub> of 0.80 under the same conditions, supporting, though not by itself proving beyond doubt, that the unknown pigment shares the identity of the reference standard; a third spot on the same plate migrating only 3.2 centimetres, giving an R<sub>f</sub> of 0.40, is clearly a distinct, more polar compound retained more strongly by the stationary phase.

**Table 9.1 Comparison of Chromatographic Techniques**

| Technique                 | Stationary Phase              | Mobile Phase                  | Typical Application                     |
|---------------------------|-------------------------------|-------------------------------|-----------------------------------------|
| Paper chromatography      | Cellulose paper               | Liquid solvent                | Separation of pigments, amino acids     |
| Thin-layer chromatography | Silica gel or alumina layer   | Liquid solvent                | Rapid qualitative separation            |
| Column chromatography     | Packed adsorbent in a column  | Liquid solvent                | Preparative-scale separation            |
| Gas chromatography        | Coated capillary column       | Inert carrier gas             | Volatile compound analysis              |
| HPLC                      | Packed column, fine particles | Liquid solvent under pressure | High-resolution quantitative separation |

**Figure 9.1 Principle of Chromatographic Separation****QUICK RECAP**

- Chromatographic separation depends on differential affinity of components for stationary and mobile phases.
- The  $R_f$  value expresses migration distance relative to the solvent front in paper and thin-layer chromatography.
- Gas chromatography suits volatile compounds; HPLC suits non-volatile compounds requiring high resolution.
- Column chromatography is used at preparative scale to isolate a purified fraction.

**COMMON ERRORS TO AVOID**

- Overloading the stationary phase with excess sample, causing streaking rather than discrete bands.
- Allowing the solvent front to run off the plate before development is complete.
- Selecting an unsuitable mobile phase polarity for the compound class under study.

**Review Questions**

[UG] 1. Define the retardation factor ( $R_f$  value) and state how it is calculated.

[UG] 2. Differentiate paper chromatography from thin-layer chromatography.

[PG] 3. Explain why gas chromatography is restricted to volatile or volatilizable compounds.

[PG] 4. Compare HPLC with conventional column chromatography in terms of resolution and operating pressure.

## Chapter 10. Electrophoretic Techniques

*Learning focus: PG core*

### Principle

Electrophoresis separates charged molecules by their differential migration through a gel matrix under an applied electric field. Migration rate depends on the net charge, size and shape of the molecule and on the pore size of the gel; in the specific case of SDS-PAGE, described below, migration depends on molecular size alone.

### Agarose Gel Electrophoresis

Agarose gel electrophoresis separates DNA and RNA fragments, which carry a uniform negative charge from their phosphate backbone, so that migration through the agarose pores depends primarily on fragment size; smaller fragments migrate faster than larger ones. Fragments are visualized under ultraviolet light after staining with an intercalating dye such as ethidium bromide, and sized by comparison with a co-run molecular weight marker.

### Gel Casting and Sample Loading

A polyacrylamide gel is cast between two glass plates as two discrete layers, a lower resolving gel of higher acrylamide concentration and a shallow upper stacking gel of low concentration and lower pH, the latter concentrating each protein sample into a sharp, narrow band before it enters the resolving gel and encounters the pore size that will separate it from its neighbours. Uneven or bubble-contaminated gel polymerization, or uneven loading volume between wells, is a frequent avoidable cause of a distorted or "smiling" band pattern that complicates accurate size comparison.

### PAGE, SDS-PAGE and Isoelectric Focusing

Native polyacrylamide gel electrophoresis (PAGE) separates proteins in their folded, native state by a combination of charge, size and shape. Treatment with sodium dodecyl sulphate (SDS), an anionic detergent that coats the polypeptide chain in proportion to its length and denatures its secondary and tertiary structure, converts SDS-PAGE into a method that separates proteins by molecular weight alone, permitting estimation of subunit size by comparison with standard proteins of known weight. Isoelectric focusing instead separates proteins along a stable pH gradient according to their isoelectric point, the pH at which a protein carries no net charge; combining isoelectric focusing with SDS-PAGE in two dimensions resolves complex protein mixtures by both charge and size.



## Blotting Techniques

Blotting transfers macromolecules resolved by electrophoresis onto a membrane for specific detection. Southern blotting detects a specific DNA sequence using a labelled complementary probe; Northern blotting applies the same principle to RNA; Western blotting transfers proteins and detects a specific protein using a labelled antibody, following the same indirect-detection logic described for immunohistochemistry in Chapter 5.

## Estimating Molecular Weight from a Gel

Molecular weight of an unknown protein or DNA fragment is estimated by plotting the logarithm of the known standard sizes against the distance each has migrated, producing an approximately linear calibration line within the resolving range of the gel; the unknown's migration distance is then read against this line rather than compared to the standards by eye alone, since visual comparison is considerably less accurate than the plotted calibration, particularly toward the extremes of the gel where migration distance changes only slightly with size.


### Worked Example: Estimating Size from a Standard Curve

A set of protein standards of known molecular weight, 116, 66, 45, 35 and 25 kilodaltons, is run alongside an unknown sample on the same SDS-PAGE gel. Plotting the logarithm of each standard's molecular weight against the distance it migrated produces an approximately straight line across this range; the unknown band, migrating to a position between the 45 and 35 kilodalton standards and closer to the 35 kilodalton marker, is read from the fitted line as approximately 38 kilodaltons. Because the relationship is only approximately linear and degrades toward the extremes of the gel, an unknown migrating close to either the largest or the smallest standard is reported as a size estimate accompanied by that caveat, rather than as an exact, high-confidence value.

**Table 10.1 Electrophoretic Techniques and Typical Matrices**

| Technique                   | Matrix             | Molecule Separated | Typical Application                   |
|-----------------------------|--------------------|--------------------|---------------------------------------|
| Agarose gel electrophoresis | Agarose gel        | DNA, RNA fragments | Nucleic acid sizing                   |
| Native PAGE                 | Polyacrylamide gel | Native proteins    | Protein separation by size and charge |

| Technique            | Matrix                          | Molecule Separated | Typical Application                    |
|----------------------|---------------------------------|--------------------|----------------------------------------|
| SDS-PAGE             | Polyacrylamide gel with SDS     | Denatured proteins | Protein molecular weight determination |
| Isoelectric focusing | pH-gradient gel                 | Proteins           | Separation by isoelectric point        |
| Western blotting     | Membrane transfer of a PAGE gel | Specific proteins  | Protein identification with antibodies |



**Figure 10.1 Steps of SDS-PAGE Analysis**

### QUICK RECAP

- Migration rate in agarose gel electrophoresis depends chiefly on nucleic acid fragment size.
- SDS coats protein in proportion to length, converting SDS-PAGE into a size-based separation.
- Isoelectric focusing separates proteins by isoelectric point along a pH gradient.
- Southern, Northern and Western blotting detect specific DNA, RNA and protein targets respectively.

### COMMON ERRORS TO AVOID

- Overheating the gel during electrophoresis by running at excessive voltage.
- Omitting a molecular weight marker, preventing accurate sizing of the resolved bands.
- Incomplete transfer during blotting, leaving target molecules within the gel rather than on the membrane.

**Review Questions**

**[UG]** 1. Explain why DNA fragments migrate toward the positive electrode during agarose gel electrophoresis.

**[UG]** 2. State the role of SDS in SDS-PAGE.

**[PG]** 3. Differentiate Southern, Northern and Western blotting by the molecule each detects.

**[PG]** 4. Explain the principle of two-dimensional gel electrophoresis and the resolution it achieves over one-dimensional SDS-PAGE.

# Chapter 11. Radioisotope Techniques in Biology

*Learning focus: PG-weighted*

## Principle

A radioisotope is an unstable atomic form that decays with emission of detectable radiation at a fixed, characteristic rate expressed as its half-life. Substituting a radioisotope for its stable counterpart within a biological molecule allows that molecule to be traced through metabolic, developmental or binding pathways by detecting the emitted radiation, without altering the chemical behaviour of the labelled molecule.

## Decline of Routine Radioisotope Use

Radioisotope methods, once basic to nucleic acid and protein labelling, have been substantially displaced in routine laboratory practice by fluorescent and chemiluminescent alternatives that avoid the licensing, shielding and waste-disposal burden associated with radioactive material, a shift evident in the fluorescent dideoxynucleotide sequencing described in Chapter 13 and the enzyme-linked detection systems described in Chapters 5, 10 and 15. Radioisotope technique nonetheless remains the method of choice for a small number of applications, particularly whole-body and whole-organ metabolic tracing, where its sensitivity and quantitative reliability have yet to be matched by a non-radioactive alternative.

## Autoradiography and Liquid Scintillation Counting

Autoradiography places a radiolabelled specimen, such as a tissue section or a chromatography plate, in contact with a photographic emulsion or a phosphor imaging plate; radioactive decay exposes the emulsion at the exact location of the label, and subsequent development yields an image showing the spatial distribution of the labelled molecule within the specimen. Liquid scintillation counting instead quantifies total radioactivity in a liquid sample: the sample is mixed with a scintillation fluid whose molecules emit a flash of light on absorbing energy from radioactive decay, and a photomultiplier tube counts these flashes to give a quantitative measure of radioactivity.

## Radiation Safety and Waste Disposal

- Handle radioactive material only in a designated, monitored work area, using shielding appropriate to the radiation type emitted.
- Wear a personal dosimeter and monitor hands and work surfaces regularly for contamination.
- Segregate radioactive waste by isotope and half-life, storing it for decay or disposal according to institutional and national regulatory requirements.

- Maintain a signed record of receipt, use and disposal for every radioactive stock.

### Choice of Isotope for a Labelling Study


Isotope selection weighs the energy and type of radiation emitted, the half-life relative to the planned duration of the experiment, and the chemical position at which the isotope can be incorporated into the molecule of interest. A short-lived, low-energy emitter such as phosphorus-32 suits a nucleic acid study completed within days, since it delivers a strong signal quickly and decays to negligible activity soon after, simplifying eventual waste disposal; a long-lived emitter such as carbon-14 suits a metabolic tracing study intended to follow a labelled compound through a slow physiological pathway over weeks.

### Worked Example: Residual Activity After Decay

A stock of phosphorus-32, with a half-life of 14.3 days, begins at an activity of 100 units on the day of receipt. After one half-life, 14.3 days, activity falls to 50 units; after two half-lives, 28.6 days, activity falls to 25 units; after four half-lives, roughly 57 days, activity has fallen below 7 units, generally considered low enough for many institutions to permit disposal as ordinary waste, subject always to the specific regulatory threshold and monitoring requirement in force locally. Planning a labelling experiment around this decay curve, rather than around the nominal activity printed on the stock's original label, ensures the working solution used on the day of the experiment actually delivers the intended signal strength.

**Table 11.1 Common Radioisotopes Used in Biological Tracing**

| Isotope              | Approximate Half-life | Typical Application             |
|----------------------|-----------------------|---------------------------------|
| Carbon-14            | 5,730 years           | Metabolic pathway tracing       |
| Phosphorus-32        | 14.3 days             | Nucleic acid labelling          |
| Iodine-125           | 59.4 days             | Protein and hormone labelling   |
| Hydrogen-3 (Tritium) | 12.3 years            | Autoradiographic cell labelling |
| Sulphur-35           | 87.5 days             | Protein synthesis studies       |



**Figure 11.1 Autoradiography Procedure**

### QUICK RECAP

- A radioisotope traces a biological molecule without altering its chemical behaviour.
- Autoradiography localizes radioactivity spatially; liquid scintillation counting quantifies total radioactivity.
- Isotope choice depends on the half-life and the type of radiation suited to the study.
- Radioactive waste must be segregated, recorded and disposed of according to regulatory requirement.

### COMMON ERRORS TO AVOID

- Selecting an isotope of unnecessarily long half-life for a short-duration experiment, complicating waste disposal.
- Working with unshielded stock radioactive material outside a designated area.
- Failing to maintain a usage and disposal record for a radioactive stock.

### Review Questions

**[UG]** 1. Define half-life and state its relevance to the choice of a radioisotope for a biological study.

**[UG]** 2. Differentiate autoradiography from liquid scintillation counting by the information each provides.

**[PG]** 3. Justify the use of phosphorus-32 over carbon-14 for the labelling of a nucleic acid probe.

**[PG]** 4. Outline the safety and record-keeping requirements for handling radioactive material in a research laboratory.

# Chapter 12. Nucleic Acid Extraction and PCR-based Techniques

## PART D · MOLECULAR BIOLOGY AND BIOINFORMATICS

*DNA-level tools basic to modern taxonomy, genetics and diagnostics*

**Learning focus:** PG core; UG conceptual

### Principle of Nucleic Acid Extraction

Extraction of DNA or RNA from animal tissue proceeds through three general stages: lysis of the cell and nuclear membrane to release nucleic acid, removal of protein and other contaminants, and precipitation or column purification of the nucleic acid in a form suitable for downstream use. Lysis is typically achieved with a detergent and a proteolytic enzyme; protein is removed by organic extraction, salting out, or binding of nucleic acid to a silica membrane under chaotropic salt conditions, followed by washing and elution.

### Quantification and Quality Assessment

Extracted nucleic acid is quantified spectrophotometrically by absorbance at 260 nanometres, with the ratio of absorbance at 260 to 280 nanometres used to assess protein contamination as described in Chapter 8, and confirmed structurally by agarose gel electrophoresis, on which intact genomic DNA appears as a single high molecular weight band and degraded DNA appears as a smear.

### Common Causes of Extraction Failure

An extraction yielding little or degraded nucleic acid usually traces to one of a small number of causes: prolonged storage of tissue at ambient temperature before lysis, allowing endogenous nuclease activity to degrade the sample; incomplete removal of a polymerase inhibitor such as residual ethanol, phenol or a polysaccharide co-extracted from certain invertebrate or plant-

associated tissue; or mechanical shearing of high molecular weight DNA during overly vigorous pipetting or vortexing. Because these failure modes are not visually distinguishable from a genuinely low starting quantity, a spectrophotometric and gel-based quality check, described below, is performed before any extract is committed to a downstream molecular procedure.

### **Principle of the Polymerase Chain Reaction**

The polymerase chain reaction (PCR) amplifies a specific DNA segment exponentially through repeated thermal cycling. Each cycle comprises three steps: denaturation, in which the double-stranded template separates into single strands at high temperature; annealing, in which two short synthetic primers bind to complementary sequences flanking the target region at a lower temperature; and extension, in which a heat-stable DNA polymerase, most commonly Taq polymerase, synthesizes a new complementary strand from each primer. Because the products of one cycle serve as templates for the next, the target sequence doubles with every cycle, yielding a large quantity of amplified product after twenty-five to thirty-five cycles.

### **Variants of PCR**

- Reverse transcription PCR (RT-PCR) first converts an RNA template into complementary DNA using the enzyme reverse transcriptase, permitting amplification of an expressed gene sequence.
- Real-time or quantitative PCR (qPCR) monitors amplification as it proceeds using a fluorescent reporter, allowing quantification of the starting template amount rather than end-point detection alone.
- Randomly amplified polymorphic DNA (RAPD) uses short, arbitrary primers to generate a strain- or population-specific band pattern useful in genetic fingerprinting.



- Restriction fragment length polymorphism (RFLP) analysis digests amplified or genomic DNA with a restriction enzyme and compares the resulting fragment pattern between individuals or species.

### **Primer Design**

An effective primer pair is specific to the target sequence, free of self-complementarity that would cause primer-dimer formation, and matched in melting temperature between the forward and reverse primer. Primer length of eighteen to twenty-five nucleotides and a guanine-cytosine content between forty and sixty percent are typical starting parameters, refined using dedicated primer-design software prior to synthesis.

### **Optimizing Annealing Temperature**

The annealing temperature is generally set a few degrees below the calculated melting temperature of the primer pair and is the single parameter most often adjusted when a PCR fails to amplify cleanly. An annealing temperature set too low permits primers to bind at partially mismatched sites, producing multiple non-specific products visible as extra bands on the confirmatory agarose gel; an annealing temperature set too high reduces primer binding efficiency at the intended site and can suppress amplification altogether. A gradient PCR, in which a single reaction is run simultaneously across a range of annealing temperatures on the same thermal cycler, is the standard method for empirically determining the optimal value for a new primer pair.


### **Worked Example: Designing a Barcode PCR Protocol**

A researcher aims to amplify the cytochrome c oxidase subunit I (COI) barcode region, approximately 650 base pairs, from a set of freshly preserved insect specimens. Using the universal LCO1490 and HCO2198 primer pair at an annealing temperature of 51 degrees Celsius, an extension time of one minute at 72 degrees Celsius, and 35 cycles, the researcher obtains a strong single band at the expected size on the confirmatory agarose gel for most

specimens, but no product at all for several specimens preserved in the field several months earlier. Because degraded template DNA amplifies poorly across a long target, the researcher repeats the failed reactions using a shorter, internal primer pair targeting only 250 base pairs of the same gene region, successfully recovering usable sequence from the degraded samples, illustrating the general principle that primer and amplicon design should be matched to template quality rather than fixed regardless of sample condition.

**Table 12.1 Components of a Standard PCR Reaction Mixture**

| Component                             | Function                                  |
|---------------------------------------|-------------------------------------------|
| Template DNA                          | Source of the target sequence             |
| Forward and reverse primers           | Define the region to be amplified         |
| Taq DNA polymerase                    | Catalyses synthesis of the new DNA strand |
| Deoxynucleotide triphosphates (dNTPs) | Building blocks for the new DNA strand    |
| Magnesium chloride                    | Cofactor required for polymerase activity |
| Reaction buffer                       | Maintains optimal pH and ionic conditions |



**Figure 12.1 Three-step PCR Thermal Cycle**

**QUICK RECAP**

- Nucleic acid extraction proceeds through lysis, contaminant removal and purification.
- PCR amplifies a target sequence exponentially through cycles of denaturation, annealing and extension.
- RT-PCR amplifies from an RNA template; qPCR quantifies template amount during amplification.
- Effective primer design balances specificity, length, guanine-cytosine content and matched melting temperature.

**COMMON ERRORS TO AVOID**

- Contaminating a DNA extraction with nuclease activity, degrading the sample before purification is complete.
- Setting an annealing temperature too low, causing non-specific primer binding and spurious products.
- Omitting a no-template control, making laboratory contamination indistinguishable from a true positive result.

**Review Questions**

- [UG] 1. List the three steps of a single PCR thermal cycle and state the purpose of each.
- [UG] 2. State the significance of the 260 to 280 nanometre absorbance ratio for extracted DNA.
- [PG] 3. Differentiate RAPD from RFLP as methods of genetic fingerprinting.
- [PG] 4. Discuss the parameters considered in the design of an effective PCR primer pair.

# CHAPTER 13. RECOMBINANT DNA TECHNOLOGY AND GENE ANALYSIS

*Learning focus: PG core*

## Principle

Recombinant DNA technology combines DNA from two different sources into a single molecule, most commonly by inserting a gene of interest into a carrier molecule, or vector, capable of independent replication within a host cell. The technology depends on two classes of enzyme: a restriction endonuclease, which cuts DNA at a specific recognition sequence, and DNA ligase, which joins compatible DNA ends to form a continuous molecule.

## Restriction Enzymes and Vectors

A restriction enzyme recognizes a short, usually palindromic, sequence of four to eight base pairs and cleaves the DNA backbone at or near that sequence, often leaving single-stranded overhangs termed sticky ends that facilitate subsequent ligation. A cloning vector, typically a bacterial plasmid, carries an origin of replication, a selectable marker such as an antibiotic resistance gene, and a cloning site into which the digested gene of interest is inserted using the same restriction enzyme, ensuring compatible ends.

## Preventing Vector Self-ligation

A linearized vector can rejoin its own two ends during the ligation step without incorporating the intended insert, producing a background of non-recombinant colonies that complicates screening. Treating the cut vector with alkaline phosphatase, which removes the terminal phosphate group required for ligation, prevents this self-ligation while leaving the insert, which retains its own terminal phosphate, free to ligate normally, substantially enriching the proportion of true recombinant colonies recovered after transformation.

## Steps in Recombinant DNA Cloning

1. Isolate the gene of interest, either from genomic DNA or as a PCR product.
2. Digest both the gene of interest and the vector with the same restriction enzyme.
3. Ligate the digested gene into the vector using DNA ligase.
4. Transform the recombinant vector into a suitable host cell, typically *Escherichia coli*.
5. Select transformed colonies on a medium containing the appropriate selective agent.

6. Screen and confirm recombinant colonies by restriction digestion or sequencing.

### DNA Sequencing and Genomic Libraries

Sanger sequencing determines nucleotide sequence through controlled synthesis of complementary strands terminated at each base position by a fluorescently labelled dideoxynucleotide, with the resulting fragments resolved by size using capillary electrophoresis. Next-generation sequencing (NGS) platforms instead sequence millions of short DNA fragments in parallel, giving far greater throughput at lower per-base cost and enabling whole-genome and metagenomic studies not feasible with Sanger sequencing alone. A genomic library represents the entire genome of an organism as a collection of cloned fragments held within a vector population, while a complementary DNA (cDNA) library represents only the genes expressed in a given tissue at a given time, prepared from messenger RNA through reverse transcription.

### Applications in Zoological Research

Recombinant DNA technology underlies the production of recombinant proteins, such as a specific hormone or enzyme, for use as an assay standard; the construction of a gene probe used in fluorescence in-situ hybridization as described in Chapter 5; and the generation of a genetically modified model organism used to study gene function directly by observing the consequence of its disruption or overexpression.


### Case Illustration: Blue-White Screening of Recombinant Colonies

A common cloning vector carries a marker gene that, when intact, allows a transformed bacterial colony to metabolize a substrate in the growth medium into a blue product, producing a visibly blue colony. Insertion of a gene of interest into the cloning site physically disrupts this marker gene, so a colony carrying a true recombinant plasmid remains white against a background of blue, non-recombinant colonies that took up the re-circularized vector alone. A researcher screening a transformation plate selects only white colonies for further analysis, then confirms each candidate by restriction digestion of its extracted plasmid, since the blue-white distinction, though a highly effective first filter, does not by itself guarantee that the insert taken up is the correct one or is present in the intended orientation.

**Table 13.1 Common Restriction Enzymes and Recognition Sequences**

| Enzyme  | Source Organism               | Recognition Sequence |
|---------|-------------------------------|----------------------|
| EcoRI   | <i>Escherichia coli</i>       | GAATTC               |
| HindIII | <i>Haemophilus influenzae</i> | AAGCTT               |

| Enzyme | Source Organism                   | Recognition Sequence |
|--------|-----------------------------------|----------------------|
| BamHI  | <i>Bacillus amyloliquefaciens</i> | GGATCC               |
| PstI   | <i>Providencia stuartii</i>       | CTGCAG               |



**Figure 13.1 Steps in Recombinant DNA Cloning**

### QUICK RECAP

- Recombinant DNA technology depends on restriction enzymes for cutting and DNA ligase for joining.
- A cloning vector supplies an origin of replication, a selectable marker and a cloning site.
- Sanger sequencing terminates synthesis at labelled bases; NGS sequences many fragments in parallel.
- A genomic library represents the whole genome; a cDNA library represents expressed genes only.

### COMMON ERRORS TO AVOID

- Using two restriction enzymes producing incompatible ends for the vector and the insert.
- Omitting a control transformation, making selection efficiency impossible to judge.
- Assuming an antibiotic-resistant colony is necessarily a true recombinant without confirmatory screening.

### Review Questions

- [UG] 1. Define restriction endonuclease and state the role of DNA ligase in cloning.
- [UG] 2. List the essential components of a bacterial cloning vector.
- [PG] 3. Compare Sanger sequencing with next-generation sequencing in principle and throughput.
- [PG] 4. Differentiate a genomic library from a complementary DNA library.

# Chapter 14. Bioinformatics Tools in Zoology

*Learning focus: PG core; UG orientation*

## Principle

Bioinformatics applies computational methods to biological sequence data, allowing a DNA or protein sequence generated in the laboratory to be compared against public reference databases for identification, functional annotation or evolutionary analysis. The approach has become basic to modern zoological taxonomy through DNA barcoding, in which a short, standardized gene region is sequenced and matched against a reference library to assign species identity.

## Basic Steps of a Bioinformatics Search

A typical bioinformatics investigation begins with a sequence generated in the laboratory, most often by the Sanger or PCR-based methods of Chapters 12 and 13, checked for quality by inspecting its chromatogram trace for ambiguous base calls before any downstream analysis is attempted. The verified sequence is then formatted according to the simple, widely used FASTA convention, a single header line beginning with the greater-than symbol followed by the sequence itself, before submission to a database search or alignment tool.

## Sequence Databases

GenBank, maintained by the National Center for Biotechnology Information (NCBI), and the European Molecular Biology Laboratory (EMBL) database are the principal international repositories of nucleotide and protein sequence data, mirrored and cross-referenced with one another. The Barcode of Life Data System (BOLD) maintains a specialized reference library of standardized barcode sequences linked to voucher specimens, supporting species-level identification from a short DNA fragment.

## Sequence Alignment and Phylogenetics

The Basic Local Alignment Search Tool (BLAST) compares a query sequence against a database and reports the closest matches ranked by similarity score, giving a rapid first indication of sequence identity. Multiple sequence alignment tools such as Clustal Omega align three or more sequences simultaneously, positioning homologous residues in common columns as a basis for further comparative or phylogenetic analysis. A phylogenetic tree, constructed from an aligned dataset using software such as MEGA, represents the inferred evolutionary relationships among the sequences through branching patterns proportional to genetic divergence.

## Representative Applications in Zoology

- Species identification from an unknown tissue sample or environmental DNA fragment.
- Confirmation of cryptic species previously indistinguishable by morphology alone.
- Reconstruction of evolutionary relationships among populations or related taxa.
- Primer design for a target gene region, and preliminary evaluation of candidate binding sites for molecular docking studies.

## Limits of Sequence-based Identification

A BLAST search reports the statistically closest matches within the database queried, not necessarily the true identity of the organism sampled, and its reliability is bounded by how completely the relevant taxonomic group is represented in that database. A high percentage identity against a poorly represented group should be treated as a provisional lead requiring confirmation by an independent line of evidence, such as morphology or a second, independent gene region, rather than as a final determination, a distinction of particular importance in biodiversity-rich but taxonomically under-sampled regions.


## Worked Example: Interpreting a BLAST Result

A COI barcode sequence obtained from an unidentified tissue sample returns a top BLAST match of 99.7 percent identity to a named species already represented by multiple accessioned records in GenBank, with the next-best match falling to 91 percent identity against a different species. Because the top match is both very high in absolute identity and clearly separated from the next-closest alternative, and because the matched region and sequence length are appropriate for the barcode marker used, the result supports confident species-level identification. A second query returning a top match of only 87 percent identity, with several other species clustered closely around a similar value, would instead be reported only to genus level or higher, and flagged as a candidate for confirmation by an independent morphological or additional-marker check before any species-level claim is published.

**Table 14.1 Commonly Used Bioinformatics Databases and Tools**

| Resource       | Primary Function                               |
|----------------|------------------------------------------------|
| GenBank (NCBI) | Repository of nucleotide and protein sequences |
| BLAST          | Sequence similarity searching                  |
| Clustal Omega  | Multiple sequence alignment                    |
| MEGA software  | Phylogenetic tree construction                 |
| BOLD Systems   | DNA barcode reference library                  |





**Figure 14.1 Workflow of Sequence-based Species Identification**

### QUICK RECAP

- Bioinformatics compares laboratory-generated sequence data against reference databases for identification and analysis.
- GenBank and EMBL are principal general sequence repositories; BOLD is specialized for DNA barcoding.
- BLAST ranks database matches by similarity; multiple sequence alignment positions homologous residues for comparison.
- A phylogenetic tree represents inferred evolutionary relationships as branching patterns.

### COMMON ERRORS TO AVOID

- Accepting a low-identity BLAST match as a confirmed species identification without corroborating evidence.
- Aligning sequences of markedly different length without appropriate trimming, distorting the alignment.
- Constructing a phylogenetic tree from an insufficiently variable gene region for the taxonomic level under study.

### Review Questions

[UG] 1. State the function of the BLAST tool in sequence analysis.

[UG] 2. Name two public sequence databases used in zoological bioinformatics.

[PG] 3. Explain the principle of DNA barcoding as a species identification method.

[PG] 4. Discuss the precautions needed before accepting a BLAST result as a confirmed species identification.

# Chapter 15. Immunological Techniques

## PART E · IMMUNOLOGY, TOXICOLOGY AND CELL CULTURE

*Assay-based techniques bridging physiology, pathology and biotechnology*

**Learning focus:** PG core; UG overview

### Principle

Immunological techniques exploit the high specificity of the antigen-antibody interaction to detect, localize or quantify a particular molecule within a biological sample. Because an antibody binds its target antigen with a specificity comparable to that of an enzyme for its substrate, the interaction underlies diagnostic tests ranging from simple agglutination to quantitative enzyme immunoassay.

### Agglutination, Precipitation and Immunodiffusion

Agglutination occurs when a multivalent antigen present on a particulate surface, such as a red blood cell or a latex bead, is cross-linked by antibody into a visible clump, forming the basis of blood grouping tests. Precipitation occurs when a soluble antigen is cross-linked by antibody into an insoluble lattice; immunodiffusion allows antigen and antibody to diffuse toward each other through a gel, forming a visible line of precipitation at the zone of optimal proportion, a pattern used to test for the identity, partial identity or non-identity of two antigen preparations.

### Polyclonal versus Monoclonal Antibodies

A polyclonal antibody preparation, raised by immunizing a host animal with the antigen of interest, contains a mixture of antibodies recognizing several distinct sites on that antigen, giving a strong but somewhat variable signal that differs between production batches. A monoclonal antibody, produced from a single cloned antibody-secreting cell line, recognizes exactly one site with high specificity and is reproducible indefinitely from batch to batch, a property that makes monoclonal reagents the preferred choice for a quantitative assay such as the sandwich ELISA described below, despite their higher cost of production.

### Enzyme-linked Immunosorbent Assay (ELISA)

ELISA quantifies an antigen or antibody using an enzyme-labelled reagent whose activity, measured colorimetrically, is proportional to the amount of target present. In the sandwich format, most widely used for antigen quantification, a capture antibody bound to the well surface binds the test antigen, an enzyme-linked detection antibody then binds a separate site on the same antigen, and addition of a substrate produces a colour proportional to bound antigen concentration, read spectrophotometrically as described in Chapter 8.

## Blood Grouping and Antibody Titre

Blood grouping applies the agglutination principle directly: red cells are mixed with antisera specific to the A, B and Rh antigens, and the pattern of agglutination observed identifies the blood group. Antibody titre, the reciprocal of the highest serum dilution still producing a detectable reaction, provides a semi-quantitative measure of antibody concentration, commonly used to follow the immune response to an experimental antigen over time.

## Constructing and Reading a Standard Curve

Quantitative ELISA results are read from a standard curve constructed by plotting the colorimetric signal produced by a dilution series of a known antigen standard against its concentration, following the same calibration logic introduced for spectrophotometry in Chapter 8. A test sample's concentration is interpolated from this curve using its measured signal, provided that signal falls within the linear portion of the curve; a sample producing a signal above the highest standard must be diluted and re-assayed rather than extrapolated beyond the calibrated range.


## Worked Example: Reading a Sandwich ELISA Plate

A sandwich ELISA standard series of 0, 5, 10, 20, 40 and 80 nanograms per millilitre of a target hormone gives absorbance readings of 0.05, 0.18, 0.34, 0.65, 1.20 and 2.05 respectively after substrate development, and plotting these values confirms an acceptable linear range up to approximately 40 nanograms per millilitre, beyond which the curve begins to plateau as capture antibody sites become saturated. A test serum sample gives an absorbance of 0.92; interpolating against the linear portion of the standard curve gives an estimated concentration of approximately 28 nanograms per millilitre. A second test sample giving an absorbance of 2.40, above the highest standard, is diluted two-fold and re-assayed rather than read directly from the plateau region of the curve, where a given absorbance no longer corresponds reliably to a single concentration value.

**Table 15.1 Types of ELISA Formats Compared**

| Format         | Principle                                            | Typical Application                  |
|----------------|------------------------------------------------------|--------------------------------------|
| Direct ELISA   | Labelled primary antibody detects antigen            | Rapid antigen screening              |
| Indirect ELISA | Labelled secondary antibody detects primary antibody | Antibody titre determination         |
| Sandwich ELISA | Antigen captured between two antibodies              | Quantification of a specific antigen |

| Format            | Principle                                           | Typical Application          |
|-------------------|-----------------------------------------------------|------------------------------|
| Competitive ELISA | Labelled and unlabelled antigen compete for binding | Detection of small molecules |



**Figure 15.1 Principle of Sandwich ELISA**

#### QUICK RECAP

- Immunological techniques depend on the specific binding of antibody to antigen.
- Agglutination involves particulate antigen; precipitation and immunodiffusion involve soluble antigen.
- Sandwich ELISA quantifies antigen using paired capture and detection antibodies.
- Antibody titre is the reciprocal of the highest reactive serum dilution.

**COMMON ERRORS TO AVOID**

- Omitting a blocking step in ELISA, allowing non-specific binding and elevated background signal.
- Interpreting weak agglutination as a negative result without adequate mixing time.
- Comparing antibody titres from assays run on different days without a shared reference standard.

**Review Questions**

- [UG] 1. Differentiate agglutination from precipitation as immunological reactions.
- [UG] 2. Describe the principle of blood grouping by the agglutination method.
- [PG] 3. Explain the sequence of binding events in a sandwich ELISA.
- [PG] 4. Define antibody titre and describe how it is determined.

# Chapter 16. Toxicological and Bioassay Techniques

*Learning focus: PG core; UG overview*

## Principle

Toxicological bioassay exposes a test organism or biological system to graded concentrations of a substance and measures a defined biological endpoint, allowing the toxicity of the substance to be expressed as a quantitative value comparable across studies. The endpoint measured, whether mortality, mutation frequency or DNA damage, determines the specific assay selected.

## LC50 and LD50 Determination

The median lethal concentration (LC50) or median lethal dose (LD50) is the concentration or dose of a substance causing death in fifty percent of a test population within a defined exposure period, commonly ninety-six hours for aquatic organisms. A preliminary range-finding trial establishes an approximate toxic range, followed by a definitive test using a closer series of concentrations bracketing the expected LC50 or LD50; mortality at each concentration is recorded and the LC50 or LD50 value is calculated from the resulting dose-response relationship, typically using probit analysis.

## Acute versus Chronic Toxicity Testing

An acute toxicity test, of which the LC50/LD50 determination is the principal example, measures the response of a test population to a single, relatively high exposure over a short, standardized period, and is primarily used to rank and classify substances by hazard. A chronic toxicity test instead exposes the test population to lower, environmentally realistic concentrations over a substantial fraction of the organism's lifespan, measuring sub-lethal endpoints such as growth rate, reproductive output or the enzyme biomarkers described below; chronic data are generally regarded as more directly relevant to real-world ecological risk than acute LC50/LD50 values alone.

## Genotoxicity Assays

The Ames test uses histidine-dependent bacterial strains to detect mutagenic activity: a test substance capable of reversing the histidine mutation allows bacterial colonies to grow on histidine-free medium, and the number of revertant colonies indicates mutagenic potency. The comet assay embeds individual cells in a thin agarose layer and subjects them to electrophoresis; damaged DNA migrates from the nucleus to form a tail resembling a comet, and tail length or

intensity indicates the degree of DNA strand breakage. The micronucleus assay scores the frequency of small, additional nuclear bodies, or micronuclei, arising from chromosome fragments or whole chromosomes not incorporated into the main nucleus during cell division, providing a measure of chromosomal damage.

### Enzyme-based Biomarkers of Toxicity

Sub-lethal exposure to a toxicant frequently alters the activity of specific enzymes before any overt sign of toxicity appears; acetylcholinesterase inhibition, for example, is a widely used biomarker of exposure to organophosphate and carbamate pesticides, measured spectrophotometrically in tissue homogenate.

### Selecting a Test Organism

A standardized aquatic toxicity test conventionally uses a sensitive, widely available and well-characterized indicator species, such as a particular daphnid, fish or amphibian larva specified in national or international test guidelines, precisely so that results obtained in different laboratories remain comparable. Substituting a locally available but non-standard species without independent validation of its sensitivity threshold undermines this comparability, even where the substitution is otherwise ecologically reasonable for the site under study.


### Worked Example: Estimating LC50 by Graphical Interpolation

A ninety-six-hour acute toxicity test on a fish species exposed to a candidate effluent gives zero percent mortality at 2 milligrams per litre, 20 percent mortality at 4 milligrams per litre, 55 percent mortality at 8 milligrams per litre, 85 percent mortality at 16 milligrams per litre and 100 percent mortality at 32 milligrams per litre. Plotting mortality against the logarithm of concentration and reading across from the fifty percent mortality mark to the fitted line gives an approximate LC50 of 7 to 8 milligrams per litre, a value that would ordinarily be refined by formal probit analysis rather than read by eye alone, but which illustrates the underlying logic: LC50 is interpolated from the dose-response relationship rather than read directly from any single tested concentration, since the concentration series is rarely designed to include the exact fifty percent point.

**Table 16.1 Toxicological Bioassay Endpoints Compared**

| Assay          | Endpoint Measured                                  | Typical Application     |
|----------------|----------------------------------------------------|-------------------------|
| LC50/LD50 test | Concentration or dose causing 50 percent mortality | Acute toxicity ranking  |
| Ames test      | Reverse mutation frequency in bacteria             | Mutagenicity screening  |
| Comet assay    | DNA strand breakage in single cells                | Genotoxicity assessment |

| Assay              | Endpoint Measured                 | Typical Application           |
|--------------------|-----------------------------------|-------------------------------|
| Micronucleus assay | Frequency of micronucleated cells | Chromosomal damage assessment |



**Figure 16.1 General Procedure for LC50/LD50 Determination**

### QUICK RECAP

- LC50 and LD50 express acute toxicity as the concentration or dose lethal to half a test population.
- The Ames test screens for mutagenicity using a bacterial reversion system.
- The comet assay and micronucleus assay measure DNA strand breakage and chromosomal damage respectively.
- Enzyme biomarkers such as acetylcholinesterase inhibition detect sub-lethal exposure before overt toxicity.

### COMMON ERRORS TO AVOID

- Omitting the preliminary range-finding trial, resulting in a definitive test that brackets the wrong concentration range.
- Using an exposure period inconsistent with the standard protocol for the species and endpoint under study.
- Failing to include a solvent control when the test substance requires a carrier solvent.

### Review Questions

[UG] 1. Define LC50 and state the exposure period conventionally used for aquatic toxicity testing.

[UG] 2. State the principle of the Ames test.

[PG] 3. Differentiate the comet assay from the micronucleus assay in the type of DNA damage each detects.

[PG] 4. Explain the value of an enzyme-based biomarker such as acetylcholinesterase inhibition in toxicological monitoring.



# Chapter 17. Animal Tissue Culture Techniques

*Learning focus: PG core*

## Principle

Animal tissue culture maintains living cells, tissue fragments or whole organ rudiments outside the body under controlled, sterile, nutritionally defined conditions. The technique allows direct experimental manipulation of cellular processes without the confounding influence of an intact organism, and underlies applications from basic cell physiology to vaccine production and toxicity screening.

## Laboratory Set-up and Sterilization

Tissue culture requires strict asepsis: work is performed within a laminar air-flow cabinet, all glassware and media are sterilized by autoclaving or filtration, and surfaces are routinely disinfected, because animal cells grow far more slowly than most microbial contaminants and a culture overrun by contamination cannot be recovered.

## Primary Culture and Established Cell Lines

A primary culture is established directly from disaggregated animal tissue and retains characteristics close to the tissue of origin, but has a finite capacity for division before senescence. An established, or continuous, cell line arises from a primary culture that has acquired the capacity for indefinite division, whether spontaneously or through deliberate transformation, and offers a standardized, reproducible system for repeated experimentation, generally at some distance from native tissue physiology.

## Culture Medium and Growth Conditions

A complete growth medium supplies a balanced salt solution, amino acids, vitamins, a carbohydrate energy source and, typically, ten percent serum providing growth factors and attachment proteins not otherwise chemically defined; antibiotics are frequently included at a low maintenance concentration as an additional safeguard against contamination, though their routine use can mask a low-grade infection until it becomes severe. Incubation is maintained at a species-appropriate temperature, commonly 37 degrees Celsius for a mammalian line, in a humidified atmosphere enriched with five percent carbon dioxide to stabilize the bicarbonate-based buffering of the medium.

## Steps in Primary Animal Cell Culture

1. Collect the tissue sample aseptically and rinse it in a balanced salt solution.

2. Disaggregate the tissue mechanically or enzymatically, commonly using trypsin or collagenase, to obtain a cell suspension.
3. Seed the cell suspension into a culture vessel containing complete growth medium.
4. Incubate at a controlled temperature and carbon dioxide concentration appropriate to the cell type.
5. Change the medium at regular intervals to replenish nutrients and remove metabolic waste.
6. Sub-culture the cells on reaching confluence, and cryopreserve stock aliquots for long-term storage.

### **Viability Assay and Cryopreservation**

The MTT assay measures cell viability and proliferation by the reduction of a tetrazolium salt to an insoluble coloured formazan product by metabolically active cells, quantified spectrophotometrically as an index of cell number or metabolic activity. Cryopreservation stores viable cells at ultra-low temperature, typically in liquid nitrogen, in the presence of a cryoprotectant such as dimethyl sulfoxide that limits ice-crystal damage during freezing and thawing.

### **Recognizing Contamination**


Bacterial or fungal contamination typically announces itself through sudden medium turbidity, an abnormal colour change in the pH indicator included in most media formulations, or visible microbial growth under low-power microscopy, and demands immediate discarding of the affected culture rather than attempted rescue, since surviving contaminants readily spread to adjacent cultures sharing the same incubator or laminar flow cabinet. Mycoplasma contamination is more insidious, producing no visible turbidity and only a subtle reduction in growth rate, and is detected only by a dedicated molecular or staining test performed at routine intervals on an ongoing culture stock.

### **Case Illustration: Deciding When to Sub-culture**

A researcher inspects a flask of adherent cells under an inverted microscope and estimates the cells cover approximately eighty-five percent of the available growth surface, with cell borders beginning to flatten against their neighbours in the most densely populated regions. Because most established lines lose their normal growth characteristics if allowed to reach full confluence and remain there, the researcher sub-cultures the flask that day rather than waiting for the culture to reach one hundred percent coverage, detaching the cells with trypsin, diluting the resulting suspension into fresh medium at a standard split ratio, and reseeding two new flasks. A parallel flask left unattended over a weekend and inspected on the following Monday shows sheets of detached, rounded cells floating in acidified, yellow-brown medium, illustrating the consequence of missing the sub-culturing window rather than monitoring confluence at the recommended interval.

**Table 17.1 Primary Culture and Established Cell Line Compared**

| Feature           | Primary Culture                     | Established Cell Line                            |
|-------------------|-------------------------------------|--------------------------------------------------|
| Source            | Directly isolated from tissue       | Derived from primary culture, often immortalized |
| Lifespan          | Limited, finite number of divisions | Extended or indefinite divisions                 |
| Genetic stability | Closer to original tissue           | May show karyotypic drift                        |
| Typical use       | Tissue-specific physiological study | Standardized, repeatable experiments             |



**Figure 17.1 Steps in Primary Animal Cell Culture**

### QUICK RECAP

- Tissue culture maintains living cells outside the body under sterile, defined conditions.
- A primary culture has finite division capacity; an established cell line divides indefinitely.
- Aseptic technique is essential because animal cells divide more slowly than most contaminants.
- The MTT assay measures viability through formazan formation by metabolically active cells.

### COMMON ERRORS TO AVOID

- Working outside the laminar flow cabinet, introducing microbial contamination.
- Allowing a culture to remain past confluence without sub-culturing, causing nutrient depletion and cell death.
- Freezing cells without an adequate cryoprotectant, causing ice-crystal damage and loss of viability on thawing.

**Review Questions**

- [UG] 1. Differentiate a primary culture from an established cell line.
- [UG] 2. State why strict asepsis is essential in animal tissue culture.
- [PG] 3. Describe the principle of the MTT assay for measuring cell viability.
- [PG] 4. Explain the role of a cryoprotectant in the cryopreservation of cultured cells.

# Chapter 18. Ecological Sampling Techniques

## PART F · ECOLOGICAL, WILDLIFE AND BEHAVIOURAL TECHNIQUES

*Field-based methods for populations, habitats and behaviour*

**Learning focus:** *Common to UG and PG study*

### Principle

Ecological sampling estimates the abundance, distribution or composition of organisms within a habitat from a representative subset of that habitat, since exhaustive enumeration is rarely feasible. The validity of any resulting estimate depends on the sampling design used, the size and number of sampling units, and the degree to which the units chosen represent the habitat as a whole.

### Quadrat and Transect Sampling

A quadrat is a defined area, commonly a one square metre frame for vegetation or intertidal sampling, placed at random or systematic intervals within the study site to record the organisms present within its boundary. A transect is a line or belt laid across an environmental gradient along which quadrats, or point observations, are taken at fixed intervals, allowing change in community composition to be related directly to the underlying gradient, such as elevation, depth or distance from a water source.

### Randomization and Replication

A sampling design gains validity from randomization, which prevents the unconscious bias of a collector who tends toward visually accessible or apparently species-rich patches, and from replication, which allows the natural variability between sampling units to be distinguished statistically from a genuine difference between sites or treatments. A single quadrat, however carefully placed, describes only itself; a defensible estimate of a site's community composition requires a number of independently placed replicates sufficient to characterize the variability actually present, a number best determined by a preliminary pilot survey rather than assumed in advance.

### Plankton and Benthos Sampling

Plankton, comprising phytoplankton and zooplankton, is sampled by towing a fine-mesh net of standard mesh size through a known volume of water, concentrating organisms for enumeration under the microscope. Benthic organisms are sampled using a grab or corer that lifts a defined volume of sediment from the bottom, which is then sieved to retrieve macroinvertebrates for identification and counting.

## Physico-chemical Analysis and Biodiversity Indices

Physico-chemical parameters of the habitat, including temperature, pH, dissolved oxygen and salinity for aquatic sites, or soil moisture and organic content for terrestrial sites, are recorded alongside biological sampling because these parameters strongly influence species distribution and must be reported together with abundance data. Biodiversity indices condense the resulting species and abundance data into a single comparative value: the Shannon-Wiener index incorporates both species richness and evenness of abundance, while the Simpson index expresses the probability that two individuals drawn at random from the sample belong to different species.

### From Raw Counts to Comparative Indices

Raw abundance counts recorded across quadrats or transects are first tabulated as a species-by-site matrix before any comparative index is calculated, since the Shannon-Wiener and Simpson indices summarized in this chapter both operate on the proportional abundance of each species within a defined sample rather than on absolute counts alone. Two sites of markedly different total abundance can nonetheless return a similar diversity index value if the relative proportions among their constituent species are similar, a distinction worth stating explicitly to students who otherwise expect diversity and abundance to track one another directly.


### Worked Example: Calculating the Shannon-Wiener Index

A quadrat survey records three species present in the following proportions of total individuals counted: species A at 0.50, species B at 0.30 and species C at 0.20. The Shannon-Wiener index is calculated as the negative sum, across all species, of each proportion multiplied by its own natural logarithm: for this sample, the contributions are approximately 0.347, 0.361 and 0.322 respectively, summing to an index value of approximately 1.03. A second site with the same three species present in exactly equal proportions of 0.333 each returns a higher index value of approximately 1.10, despite having identical species richness, because the Shannon-Wiener index rewards evenness of abundance in addition to the mere number of species present, precisely the property that a simple species count would fail to capture.

**Table 18.1 Common Ecological Sampling Methods and Their Targets**

| Method                | Habitat                 | Organism Group                               |
|-----------------------|-------------------------|----------------------------------------------|
| Quadrat sampling      | Terrestrial, intertidal | Vegetation, sessile invertebrates            |
| Belt transect         | Terrestrial, aquatic    | Distribution along an environmental gradient |
| Plankton net sampling | Open water              | Phytoplankton, zooplankton                   |

| Method                    | Habitat                  | Organism Group             |
|---------------------------|--------------------------|----------------------------|
| Sediment sampling<br>grab | Benthic zone             | Benthic macroinvertebrates |
| Pitfall trapping          | Terrestrial soil surface | Ground-dwelling arthropods |



Numbered quadrats placed at fixed intervals along the transect line

**Figure 18.1 Quadrat Sampling Layout within a Study Transect**

### QUICK RECAP

- Ecological sampling estimates population and community parameters from a representative subset of the habitat.
- Quadrat sampling records organisms within a defined area; transect sampling relates composition to an environmental gradient.
- Plankton is sampled by net tow; benthos is sampled by grab or corer followed by sieving.
- The Shannon-Wiener and Simpson indices summarize species richness and evenness in a single comparative value.

### COMMON ERRORS TO AVOID

- Placing quadrats non-randomly by unconscious preference for visually rich patches, biasing the estimate.
- Using too few sampling units to characterize the variability present in the habitat.
- Recording biological data without accompanying physico-chemical measurements needed for interpretation.

### Review Questions

[UG] 1. Define quadrat sampling and state one habitat in which it is commonly applied.

[UG] 2. Differentiate plankton sampling from benthos sampling in method and target organisms.

- [PG]** 3. Explain what the Shannon-Wiener index measures beyond simple species richness.
- [PG]** 4. Discuss the design considerations affecting the reliability of a transect-based ecological survey.



# Chapter 19. Wildlife Census and Population Estimation Techniques

*Learning focus: PG-weighted; UG overview*

## Principle

Wildlife census techniques estimate the size, density or distribution of a free-ranging animal population without requiring enumeration of every individual, an approach necessary because direct total counts are rarely practical for mobile, cryptic or wide-ranging species. Method selection depends on the mobility, detectability and identifiability of the species under study.

## Direct and Indirect Count Methods

Direct count methods record animals actually observed, whether by ground survey, vantage-point count or aerial survey, and are best suited to conspicuous species in open habitat. Indirect methods infer presence and abundance from signs left by the animal, including tracks, scat, burrows, calls or camera-trap photographs, and are preferred for cryptic, nocturnal or low-density species in closed habitat.

## Marking Methods and Their Selection

A marking method must be visible or detectable throughout the study period, must not alter the marked individual's behaviour or survival relative to unmarked individuals, and must be appropriate to the biology of the species concerned. Options range from a numbered leg band or ear tag for a vertebrate, through a fin clip or elastomer tag for a fish, to a fluorescent dust or paint mark for many invertebrates; a method causing even a slight reduction in survival or an increased conspicuousness to predators biases the resulting population estimate and must be piloted before adoption in a full study.

## Mark-recapture and Line Transect Methods

The mark-recapture method captures, marks and releases a sample of individuals, then captures a second independent sample after a defined interval and records the proportion of marked individuals within it; the Lincoln-Petersen estimator uses this proportion to calculate total population size, on the assumption that marked and unmarked individuals mix freely and are equally likely to be recaptured. Line transect distance sampling instead records the perpendicular distance from a fixed transect line to each animal or animal sign detected, and uses the resulting distribution of detection distances to estimate the effective area surveyed and, from it, population density.

## Camera Trapping, Telemetry and Sign-based Census

Camera trapping deploys motion-triggered cameras at fixed stations to photograph passing animals; for species with individually distinctive markings, such as tiger stripe patterns, repeated photographic capture of identified individuals over multiple stations supports a capture-recapture population estimate without physical handling. Radio-telemetry and satellite tracking attach a transmitter to a captured individual, allowing its location to be recorded remotely over an extended period, providing data on home range, movement pattern and habitat use rather than population size directly. Pugmark and scat-based census methods, long used for large carnivores in the Indian subcontinent, identify individuals from track dimensions or estimate relative abundance from sign density along a standardized survey route.

### Population Density from an Estimate

A population estimate obtained by mark-recapture or a related method describes the number of individuals present within the effectively sampled area, and must be divided by that area to yield density, a step frequently overlooked by students who report an estimate as though it applied automatically to the entire study region. Correctly defining the effectively sampled area, accounting for animal movement across the boundary of a trapping grid during the study period, is itself a well-recognized methodological difficulty addressed by dedicated spatial capture-recapture models in advanced practice.


### Worked Example: Applying the Lincoln-Petersen Estimator

A first trapping session captures and marks 40 individuals of a small mammal species, which are then released back into the study area. A second trapping session conducted one week later captures 50 individuals, of which 10 carry a mark from the first session. The Lincoln-Petersen estimator calculates total population size as the first sample size multiplied by the second sample size, divided by the number of marked recaptures: 40 multiplied by 50, divided by 10, giving an estimated population of 200 individuals. Had only 4 marked individuals been recaptured instead of 10, the same formula would give an estimate of 500, illustrating how sensitive the estimator is to the recapture count and why an adequately large second sample is essential to a reasonably precise estimate.

**Table 19.1 Wildlife Population Estimation Methods Compared**

| Method                          | Data Required                           | Suitable Species                          |
|---------------------------------|-----------------------------------------|-------------------------------------------|
| Mark-recapture                  | Marked and recaptured individual counts | Mobile, individually identifiable animals |
| Line transect distance sampling | Perpendicular sighting distances        | Visually detectable animals               |
| Camera trap capture-recapture   | Photographic identification records     | Individually patterned mammals            |

| Method                 | Data Required                         | Suitable Species |
|------------------------|---------------------------------------|------------------|
| Pugmark / track census | Track measurements along fixed routes | Large carnivores |



**Figure 19.1 Mark-recapture Protocol (Lincoln-Petersen Method)**

### QUICK RECAP

- Direct methods count animals observed; indirect methods infer presence from signs.
- The Lincoln-Petersen estimator calculates population size from the proportion of marked recaptures.
- Line transect distance sampling estimates density from the distribution of detection distances.
- Radio-telemetry provides movement and home-range data rather than a direct population estimate.

### COMMON ERRORS TO AVOID

- Assuming free mixing of marked and unmarked individuals when territoriality or trap-shyness violates that assumption.
- Using too short a time interval between mark and recapture samples, so marks have not yet mixed into the population.
- Misidentifying individuals from ambiguous camera-trap images, inflating the apparent population estimate.

### Review Questions

[UG] 1. State the assumption underlying the Lincoln-Petersen mark-recapture estimator.

[UG] 2. Differentiate direct from indirect wildlife census methods.

[PG] 3. Explain the data recorded in line transect distance sampling and its use in estimating density.

[PG] 4. Discuss the advantage of camera-trap capture-recapture over physical mark-recapture for elusive carnivores.

# Chapter 20. Remote Sensing, GIS and Animal Behaviour Study

*Learning focus: PG-weighted; UG orientation*

## Principle of Remote Sensing and GIS

Remote sensing acquires information about the Earth's surface from satellite or aerial sensors without direct physical contact, while a Geographic Information System (GIS) stores, analyses and displays that information, together with field-collected data, as spatially referenced layers. In wildlife ecology, the combination allows habitat characteristics to be mapped and related quantitatively to animal distribution recorded by field survey or telemetry.

## GPS and Habitat Mapping

A Global Positioning System (GPS) receiver records the geographic coordinates of a field observation, specimen locality or telemetry fix with an accuracy typically within a few metres, providing the spatial reference required to enter field data into a GIS. Common data layers used in habitat mapping include land cover classification derived from satellite imagery, a digital elevation model describing terrain and slope, and a vegetation index such as the Normalized Difference Vegetation Index (NDVI), which estimates vegetation density and health from the differential reflectance of healthy vegetation in the red and near-infrared bands.

## Species Distribution Modelling

Field GPS records of confirmed species occurrence, combined with the environmental data layers described above, can be used to build a species distribution model, a statistical relationship between recorded presence and the underlying habitat variables that is then projected across the wider landscape to predict where the species is likely to occur beyond the points actually surveyed. Such models are only as reliable as the environmental layers and the geographic spread of the occurrence records feeding them, and a model built from a geographically clustered or environmentally narrow set of records should be interpreted cautiously outside the range it was trained on.

## Principle of Ethogram-based Behaviour Study

An ethogram is a structured, exhaustive catalogue of the discrete behaviour patterns displayed by a species, each behaviour defined precisely enough that independent observers record it consistently. Once an ethogram is established, standardized sampling rules are applied to convert continuous behaviour into quantifiable data: focal-animal sampling follows one identified individual for a fixed period and records every occurrence of each defined behaviour,

while scan sampling records the behaviour of every visible individual in a group at fixed time intervals, giving a population-level rather than an individual-level picture.

### **Focal Animal Sampling Protocol**

1. Select a focal individual for observation, ensuring it can be followed for the full sampling period.
2. Define the sampling period and interval in advance of observation.
3. Record continuously every occurrence of each ethogram-defined behaviour shown by the focal individual.
4. Code the recorded behaviour according to the established ethogram categories.
5. Compile the coded data into a behavioural time budget expressing the proportion of time spent in each activity.

### **Inter-observer Reliability**

Because behavioural data depend on human judgement in a way that instrument-based measurement does not, a behavioural study should establish inter-observer reliability before data collection begins, typically by having two trained observers independently code the same recorded sequence and calculating the percentage agreement or an equivalent statistic between their records. Reliability below an accepted threshold, commonly around eighty to eighty-five percent agreement, indicates that the ethogram's behavioural definitions require further refinement before the study proceeds, rather than a problem to be addressed after data collection is already complete.

### **Bioacoustics**

Bioacoustic methods record and analyse animal sound using directional microphones and spectrographic software, allowing species identification, individual recognition and behavioural context to be inferred from call structure, a technique of particular value for cryptic or nocturnal species that are rarely seen but frequently heard.


### **Worked Example: Building a Behavioural Time Budget**

A focal-animal sampling session on a captive primate over a sixty-minute observation period records 24 minutes of feeding, 18 minutes of resting, 12 minutes of locomotion and 6 minutes of social grooming, coded continuously against a previously validated ethogram. Converting each behaviour to a percentage of the total observation period gives a time budget of 40 percent feeding, 30 percent resting, 20 percent locomotion and 10 percent grooming. Repeating this same protocol across multiple individuals and multiple days, then averaging the resulting time budgets, allows the researcher to describe a representative activity pattern for the study

population and to test statistically, using the methods of Chapter 21, whether that pattern differs between, for example, a wild and a captive population of the same species.

**Table 20.1 GIS and Remote Sensing Data Layers in Habitat Mapping**

| Data Layer                | Source                       | Application                   |
|---------------------------|------------------------------|-------------------------------|
| Land cover classification | Satellite imagery            | Habitat type delineation      |
| Digital elevation model   | Satellite or aerial survey   | Terrain and slope analysis    |
| Vegetation index (NDVI)   | Multispectral satellite data | Vegetation health and density |
| Species occurrence points | Field GPS records            | Distribution mapping          |



**Figure 20.1 Focal Animal Sampling Protocol**

### QUICK RECAP

- Remote sensing supplies habitat data; GIS integrates that data with field observations as spatial layers.
- NDVI estimates vegetation density from differential red and near-infrared reflectance.
- An ethogram is a precisely defined catalogue of behaviour patterns basic to consistent behavioural recording.
- Focal-animal sampling records one individual continuously; scan sampling records a group at fixed intervals.

### COMMON ERRORS TO AVOID

- Using an ethogram with ambiguously defined categories, reducing agreement between independent observers.
- Selecting a focal individual that repeatedly moves out of observable range during the sampling period.
- Interpreting a single NDVI image without accounting for seasonal variation in vegetation cover.

**Review Questions**

[UG] 1. Define ethogram and state its role in behavioural sampling.

[UG] 2. Differentiate focal-animal sampling from scan sampling.

[PG] 3. Explain the principle by which NDVI estimates vegetation density from satellite reflectance data.

[PG] 4. Discuss the value of bioacoustic recording for the study of cryptic or nocturnal species.

# Chapter 21. Biostatistics for Zoological Data

*Learning focus: Common to UG and PG study*

## Principle

Biostatistics applies statistical method to biological data, converting raw measurements or counts into summary values and formal tests that support or refute a stated hypothesis. Sound statistical practice begins before data collection, with a sampling design adequate to the question, and continues through data description, hypothesis testing and interpretation of the result within its biological context.

## Data Types and Presentation

Biological data are classified as qualitative, categorizing observations into non-numeric classes such as sex or species, or quantitative, expressed as numeric counts (discrete data) or measurements (continuous data). Tabulation and graphical presentation, including frequency tables, bar diagrams, histograms and scatter plots, condense raw data into an interpretable form and frequently reveal a pattern not apparent from a simple list of values.

## Measures of Central Tendency, Dispersion and Probability

The mean, median and mode describe the central value of a dataset, each suited to a different data distribution or scale. Variance and standard deviation describe the spread of values about the mean, while the standard error describes the precision of the mean itself as an estimate of the true population value. Probability theory underlies inferential statistics: a dataset approximating the normal distribution supports the parametric tests described below, while a markedly skewed or ordinal dataset generally requires a non-parametric alternative; the Poisson distribution, describing the probability of rare, independent events, is commonly relevant to count data such as the number of parasites per host.

## Null Hypothesis and Type I and Type II Error

Every test of significance evaluates a null hypothesis, typically the assertion that no true difference or association exists, against an alternative hypothesis proposing that one does. Rejecting a true null hypothesis is termed a Type I error, conventionally limited to a five percent probability by the chosen significance threshold; failing to reject a false null hypothesis, most often because the study sample was too small to detect a real but modest effect, is termed a Type II error. Statistical power, the probability of correctly detecting a true effect, is increased chiefly by increasing sample size, and should be estimated at the study design stage rather than discovered as an unwelcome explanation for a non-significant result obtained afterward.

## Tests of Significance



A test of significance evaluates whether an observed difference or association is unlikely to have arisen by chance alone, expressed as a probability value, or p-value, against a conventional threshold, usually 0.05. Student's t-test compares the means of two groups; one-way analysis of variance (ANOVA) extends the comparison to three or more groups; the chi-square test evaluates association or goodness of fit for categorical, frequency-based data.

## **Correlation, Regression and Statistical Software**

Correlation measures the strength and direction of a linear association between two continuous variables, expressed by the Pearson correlation coefficient, while regression models the quantitative relationship between a dependent variable and one or more independent variables, supporting prediction as well as description of association. Statistical software, ranging from spreadsheet functions in MS Excel to dedicated packages such as SPSS and the R programming environment, performs these calculations reliably at a scale impractical by hand and additionally supports the graphical output basic to result presentation.

## **Reporting Statistical Results**


A statistical result reported in a thesis or manuscript should state the test used, the test statistic obtained, the associated degrees of freedom where applicable, and the exact or bounded p-value, presented alongside a descriptive summary such as the mean and standard error rather than the p-value in isolation. Reporting only that a result was "significant," without these accompanying values, prevents a reader from independently judging the strength of the evidence and is inconsistent with the transparency expected of a scientific publication.

## **Worked Example: Applying Student's t-test**

A researcher compares body weight between a control group of ten rats and a treatment group of ten rats, obtaining a mean of 220 grams with a standard deviation of 15 in the control group and a mean of 205 grams with a standard deviation of 14 in the treatment group. Applying an independent-samples t-test to these two groups gives a t-statistic of approximately 2.34 with 18 degrees of freedom, corresponding to a p-value of approximately 0.031, below the conventional 0.05 threshold, so the null hypothesis of no difference between groups is rejected and the weight difference is reported as statistically significant. The researcher additionally reports the actual mean difference of 15 grams and its confidence interval alongside the p-value, since the p-value alone states only that a difference exists, not its likely biological magnitude.

**Table 21.1 Common Statistical Tests and Their Applications**

| Test                | Data Type                                       | Purpose                                 |
|---------------------|-------------------------------------------------|-----------------------------------------|
| Student's t-test    | Continuous, two groups                          | Comparison of two means                 |
| One-way ANOVA       | Continuous, three or more groups                | Comparison of multiple means            |
| Chi-square test     | Categorical / frequency data                    | Test of independence or goodness of fit |
| Pearson correlation | Continuous, paired data                         | Strength of linear association          |
| Linear regression   | Continuous, dependent and independent variables | Prediction and relationship modelling   |

**Figure 21.1 Decision Pathway for Selecting a Statistical Test****QUICK RECAP**

- Statistical design begins before data collection with an adequate sampling plan.
- Mean, median and mode summarize central tendency; variance and standard deviation summarize spread.
- The t-test, ANOVA and chi-square test address two-group, multi-group and categorical comparisons respectively.
- Correlation measures association strength; regression models and predicts a quantitative relationship.

**COMMON ERRORS TO AVOID**

- Applying a parametric test to markedly non-normal data without checking the distribution assumption.
- Treating a statistically significant result as automatically biologically important.
- Confusing correlation with a causal relationship between two variables.

**Review Questions**

[UG] 1. Differentiate the mean, median and mode as measures of central tendency.

[UG] 2. State the purpose of the chi-square test and the data type to which it applies.

[PG] 3. Explain the distinction between correlation and regression.

[PG] 4. Discuss the assumptions that must be checked before applying a parametric test of significance.

# Chapter 22. Research Methodology, Thesis Writing, Ethics and Intellectual Property

*Learning focus: PG core; UG orientation*

## **Research Design and Literature Review**

A zoological research project begins with a clearly stated problem, refined into a testable hypothesis through a systematic review of existing literature that identifies what is already known, what remains unresolved, and how the proposed study will add to that knowledge. The study design, encompassing the sampling strategy, techniques to be applied and the statistical analysis planned, should be finalized before data collection begins, since a design flaw discovered afterward is rarely correctable.

## **Referencing and Scientific Writing**

Every factual claim, method or figure drawn from another source requires accurate citation, both to credit the original author and to allow a reader to verify the claim independently; common referencing styles include the author-date format and the numbered reference format, the specific choice generally fixed by the target journal or institution. Scientific writing favours precision over elaboration: a claim is stated once, supported by evidence, and expressed in the past tense for completed work and the present tense for generally accepted fact.

## **Preparing the Manuscript for Peer Review**

Submission to a peer-reviewed journal follows the target journal's specific formatting and referencing requirements, detailed in its instructions to authors, and is generally accompanied by a cover letter stating the manuscript's novelty and its fit to the journal's scope. Reviewer comments returned after the initial submission should be addressed point by point in a structured response letter, distinguishing a change actually made to the manuscript from a reasoned justification for a suggestion not adopted, since an unaddressed reviewer comment is a common and avoidable cause of an otherwise sound manuscript's rejection.

## **Thesis Structure and Plagiarism**

A thesis or research manuscript conventionally follows the IMRaD structure: Introduction, Materials and Methods, Results, and Discussion, preceded by an abstract and followed by a reference list, as summarized in Table 22.1. Plagiarism, the presentation of another person's words, data or ideas as one's own without acknowledgement, is a serious breach of academic integrity regardless of intent, and is routinely screened for using similarity-detection software prior to submission.

## **Animal Ethics**

Research involving live vertebrate animals requires prior approval from an Institutional Animal Ethics Committee, constituted in India under the Committee for the Purpose of Control and Supervision of Experiments on Animals (CPCSEA), which reviews the proposed procedure against the principles of replacement, reduction and refinement before permitting the study to proceed. Field research on wild populations similarly requires compliance with relevant wildlife protection legislation and collection permits issued by the appropriate forest or wildlife authority.

## **Intellectual Property Rights**

Intellectual property rights protect the outputs of research with commercial or applied value: a patent protects a novel, non-obvious and industrially applicable invention for a defined term, while copyright protects the original expression of a published work, including a thesis, without requiring formal registration. A researcher developing a novel diagnostic method, bioassay protocol or biotechnological process should establish patentability, generally through prior disclosure to the institution's technology transfer office, before public disclosure of the work in a conference or publication, since prior public disclosure can itself defeat a later patent claim.

## **The Three R Principles in Animal Research**


The three R principles evaluated by an Institutional Animal Ethics Committee are replacement, the use of a non-animal or lower-sentience alternative wherever it can answer the same scientific question; reduction, the use of the minimum number of animals statistically sufficient to obtain a reliable result, informed by the power consideration introduced in Chapter 21; and refinement, the modification of a procedure to minimize pain, distress or lasting harm to the animals actually used. A research proposal that has not explicitly considered all three principles is generally returned by the committee for revision before any approval is granted.

## **Case Illustration: From Objective to Publishable Result**

A postgraduate student begins with a broad interest in the effect of a wetland pollutant on local amphibian populations, and through the literature review narrows this into the testable hypothesis that the pollutant reduces tadpole growth rate at concentrations already reported in the affected wetland. The study design specifies a laboratory bioassay following the LC50/LD50 logic of Chapter 16 at sub-lethal concentrations, growth measured at fixed intervals, and a one-way ANOVA comparing growth rate across concentration groups as set out in Chapter 21. Data collection, statistical analysis and manuscript drafting each follow directly from decisions made at the design stage, and the completed manuscript, structured according to Table 22.1, is submitted only after Institutional Animal Ethics Committee approval obtained before the study began is cited in the methods section, closing the loop from the original broad interest to a specific, defensible, publishable result.

**Table 22.1 IMRaD Structure of a Zoological Thesis or Manuscript**

| Section                   | Content                                                     |
|---------------------------|-------------------------------------------------------------|
| Introduction              | Background, rationale, objectives                           |
| Materials and Methods     | Study area, sampling, techniques used, statistical analysis |
| Results                   | Findings presented through text, tables and figures         |
| Discussion                | Interpretation of results in light of existing literature   |
| Conclusion and References | Summary of findings, cited literature                       |

**Figure 22.1 Workflow of a Zoological Research Project****QUICK RECAP**

- A testable hypothesis is refined from a stated problem through systematic literature review.
- The IMRaD structure organizes a thesis or manuscript into Introduction, Methods, Results and Discussion.
- Institutional Animal Ethics Committee approval, under CPCSEA guidelines, is mandatory before vertebrate research begins.
- A patent requires novelty and is defeated by public disclosure occurring before an application is filed.

**COMMON ERRORS TO AVOID**

- Finalizing the study design after data collection has already begun, leaving flaws that cannot be corrected.
- Paraphrasing a source closely enough to constitute plagiarism despite the absence of a direct quotation.
- Commencing vertebrate animal research before Institutional Animal Ethics Committee approval is obtained.

**Review Questions**

**[UG]** 1. List the sections of the IMRaD thesis structure in order.

**[UG]** 2. Define plagiarism and state why it is regarded as a breach of academic integrity.

**[PG]** 3. Explain the role of an Institutional Animal Ethics Committee under CPCSEA guidelines.

**[PG]** 4. Discuss why public disclosure of a novel method before filing can defeat a subsequent patent claim.

## Appendix I · Glossary of Technical Terms

Concise definitions of terms used repeatedly across this book. A term defined in detail within a specific chapter is not repeated here.

| Term                      | Definition                                                                                                        |
|---------------------------|-------------------------------------------------------------------------------------------------------------------|
| <b>Absorbance</b>         | The logarithmic measure of light attenuation by a sample, basic to spectrophotometric quantification.             |
| <b>Agglutination</b>      | Clumping of particulate antigen caused by cross-linking antibody.                                                 |
| <b>Amplicon</b>           | The DNA product generated by a polymerase chain reaction.                                                         |
| <b>Antigen retrieval</b>  | Restoration of antibody-binding ability lost during tissue fixation.                                              |
| <b>Autolysis</b>          | Self-digestion of tissue by its own enzymes after death, arrested by fixation.                                    |
| <b>Autoradiography</b>    | Localization of a radiolabelled molecule by its exposure of a photographic emulsion.                              |
| <b>Bioassay</b>           | An experimental procedure that measures a biological response to quantify the potency or toxicity of a substance. |
| <b>Biodiversity index</b> | A single numerical value summarizing species richness and evenness within a community.                            |
| <b>Blocking</b>           | Saturation of non-specific binding sites in an immunoassay before the specific reagent is applied.                |
| <b>Calibration curve</b>  | A graph relating instrument response to the concentration of known standards.                                     |
| <b>Chaotropic salt</b>    | A salt that disrupts hydrogen bonding, used to denature protein during nucleic acid purification.                 |



| Term                     | Definition                                                                                                            |
|--------------------------|-----------------------------------------------------------------------------------------------------------------------|
| <b>Colchicine</b>        | An alkaloid that arrests dividing cells at metaphase by inhibiting spindle formation.                                 |
| <b>Confluence</b>        | The state of a cell culture in which cells cover the entire available surface.                                        |
| <b>Cryoprotectant</b>    | A chemical agent that limits ice-crystal damage to cells during freezing.                                             |
| <b>Denaturation</b>      | Loss of the native structure of a protein or nucleic acid, whether by heat or chemical agent.                         |
| <b>Ethogram</b>          | A structured catalogue of the discrete behaviour patterns displayed by a species.                                     |
| <b>Fixation</b>          | Chemical or physical stabilization of tissue structure, arresting decomposition.                                      |
| <b>Formazan</b>          | The coloured, insoluble product formed from a tetrazolium salt by metabolically active cells, basic to the MTT assay. |
| <b>Genotoxicity</b>      | The capacity of an agent to damage the genetic material of a cell.                                                    |
| <b>Half-life</b>         | The time required for half of a given quantity of a radioisotope to decay.                                            |
| <b>Histochemistry</b>    | Localization of a specific chemical constituent within a tissue section by a chemically specific reaction.            |
| <b>Homogenate</b>        | A tissue sample disrupted into a uniform suspension of cellular and subcellular components.                           |
| <b>Idiogram</b>          | A diagrammatic representation of a karyotype arranged by chromosome size and morphology.                              |
| <b>Isoelectric point</b> | The pH at which a protein carries no net electrical charge.                                                           |

| Term                      | Definition                                                                                                     |
|---------------------------|----------------------------------------------------------------------------------------------------------------|
| <b>Karyotype</b>          | The ordered chromosome complement of an organism, displayed by size and morphology.                            |
| <b>Ligation</b>           | Enzymatic joining of two DNA fragments by DNA ligase.                                                          |
| <b>Mark-recapture</b>     | A population estimation method based on the proportion of marked individuals recovered in a second sample.     |
| <b>Metaphase</b>          | The stage of cell division at which chromosomes are maximally condensed and aligned for analysis.              |
| <b>Micronucleus</b>       | A small additional nuclear body formed from a chromosome fragment not incorporated into the main nucleus.      |
| <b>Monochromator</b>      | An optical component that isolates a narrow band of wavelength from a light source.                            |
| <b>Numerical aperture</b> | A measure of the light-gathering ability of a microscope objective, determining resolving power.               |
| <b>Passage</b>            | A single cycle of sub-culturing a cell line from one vessel to another.                                        |
| <b>Phylogenetic tree</b>  | A branching diagram representing inferred evolutionary relationships among sequences or taxa.                  |
| <b>Primer</b>             | A short synthetic oligonucleotide that initiates DNA synthesis at a specific site.                             |
| <b>Probit analysis</b>    | A statistical method converting a sigmoid dose-response curve into a linear form for calculating LC50 or LD50. |
| <b>Quadrat</b>            | A defined sampling area used to record organisms present within its boundary.                                  |

| Term                            | Definition                                                                                                          |
|---------------------------------|---------------------------------------------------------------------------------------------------------------------|
| <b>Reagent blank</b>            | A control sample containing all reagents except the analyte, used to zero an instrument.                            |
| <b>Recombinant</b>              | A DNA molecule formed by joining genetic material from two different sources.                                       |
| <b>Restriction endonuclease</b> | An enzyme that cleaves DNA at a specific recognition sequence.                                                      |
| <b>Retardation factor (Rf)</b>  | The ratio of the distance travelled by a compound to the distance travelled by the solvent front in chromatography. |
| <b>Sedimentation</b>            | The settling of particles under gravitational or centrifugal force.                                                 |
| <b>Titre</b>                    | The reciprocal of the highest dilution of a serum still producing a detectable reaction.                            |
| <b>Vector</b>                   | A DNA molecule, typically a plasmid, used to carry a gene of interest into a host cell.                             |

### *About Editors*



**Mr. Kishor U. More** is working as an Assistant Professor at Mudhoji College, Phaltan. He completed his B.Sc. and M.Sc. in Zoology from Savitribai Phule Pune University, Pune, and qualified the CSIR-NET (Life Science) examination. He has 10 years of teaching experience and is actively engaged in teaching and academic research. His research interests focus on entomology and environmental sciences. He has published 6 research papers in national and international peer-reviewed journals and authored one book chapter. He is the inventor of 2 patents and has actively participated in 5 national and International conferences, as well as 17 workshops. He was Member of the Board of Studies in the Department of Zoology at Tuljaram Chaturchand College, Baramati.



**Miss. Sharvari V. Shah** is working as an Assistant Teacher at Vidya Pratishthan's Dr. Cyrus Poonawalla School (CBSE), Rui, Vidyanagari, Baramati, Maharashtra. She completed her B.Sc. in Zoology, M.Sc. in Entomology, and B.Ed. from Savitribai Phule Pune University, Pune. She has 5 years of teaching experience and is actively engaged in teaching and academic research. Her research interests focus on entomology, biodiversity, and environmental sciences. She has published 4 research papers in national and international peer-reviewed journals. She is the co-inventor of two patents and has actively participated in 2 national conferences and 5 workshops. She was Member of the Board of Studies in the Department of Zoology at Tuljaram Chaturchand College, Baramati.



**Dr. Kavita Kumbhar** received her M. Sc. Degree in Zoology in 2011, B.Ed. in 2012 and in March 2024 she declared Ph.D. in Zoology (Entomology) from Shivaji University, Kolhapur, Maharashtra. She has presented 03 papers in National and International conferences. She has published 01 book and 05 research papers in a National and International Journals. She has 12 years of teaching experience at under graduate level. She has worked as a Head and assistant professor in Zoology, at Shri Shahaji Chhatrapati Mahavidhyalaya, Kolhapur. Currently she is working as an Entomologist under National Health Mission, Kolhapur.



**Miss. Shaheen M. Shaikh** is working as an Assistant Teacher at ADT's Shardabai Pawar Mahila Mahavidyalaya, Shardanagar, Maharashtra. She completed her B.Sc. and M.Sc. in Zoology from Savitribai Phule Pune University, Pune. She has 3 years of teaching experience and is actively engaged in teaching and academic development.

**MRP. Rs. 299/-**

**ISBN: 978-81-18-05443-6**



9 788118 054436